





# BMJ Open Non-operative treatment for lumbar spinal stenosis with neurogenic claudication: an updated systematic review

Carlo Ammendolia <sup>1,2</sup> Corey Hofkirchner,<sup>3</sup> Joshua Plener,<sup>3</sup> André Bussi  res <sup>4,5</sup> Michael J Schneider,<sup>6</sup> James J Young <sup>3,7</sup>, Andrea D Furlan <sup>8,9</sup> Kent Stuber,<sup>3</sup> Aksa Ahmed,<sup>2</sup> Carol Cancelliere,<sup>10</sup> Aleisha Adeboyejo,<sup>3</sup> Joseph Ornelas<sup>11</sup>

**To cite:** Ammendolia C, Hofkirchner C, Plener J, *et al.* Non-operative treatment for lumbar spinal stenosis with neurogenic claudication: an updated systematic review. *BMJ Open* 2022;**12**:e057724. doi:10.1136/bmjopen-2021-057724

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-057724>).

Received 27 September 2021  
Accepted 15 December 2021



  Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Carlo Ammendolia;  
[cammendolia@mtsina1.on.ca](mailto:cammendolia@mtsina1.on.ca)

## ABSTRACT

**Objectives** Neurogenic claudication due to lumbar spinal stenosis (LSS) is a growing health problem in older adults. We updated our previous Cochrane review (2013) to determine the effectiveness of non-operative treatment of LSS with neurogenic claudication.

**Design** A systematic review.

**Data sources** CENTRAL, MEDLINE, EMBASE, CINAHL and Index to Chiropractic Literature databases were searched and updated up to 22 July 2020.

**Eligibility criteria** We only included randomised controlled trials published in English where at least one arm provided data on non-operative treatment and included participants diagnosed with neurogenic claudication with imaging confirmed LSS.

**Data extraction and synthesis** Two independent reviewers extracted data and assessed risk of bias using the Cochrane Risk of Bias Tool 1. Grading of Recommendations Assessment, Development and Evaluation was used for evidence synthesis.

**Results** Of 15200 citations screened, 156 were assessed and 23 new trials were identified. There is moderate-quality evidence from three trials that: Manual therapy and exercise provides superior and clinically important short-term improvement in symptoms and function compared with medical care or community-based group exercise; manual therapy, education and exercise delivered using a cognitive-behavioural approach demonstrates superior and clinically important improvements in walking distance in the immediate to long term compared with self-directed home exercises and glucocorticoid plus lidocaine injection is more effective than lidocaine alone in improving statistical, but not clinically important improvements in pain and function in the short term. The remaining 20 new trials demonstrated low-quality or very low-quality evidence for all comparisons and outcomes, like the findings of our original review.

**Conclusions** There is moderate-quality evidence that a multimodal approach which includes manual therapy and exercise, with or without education, is an effective treatment and that epidural steroids are not effective for the management of LSS with neurogenic claudication. All other non-operative interventions provided insufficient quality evidence to make conclusions on their effectiveness.

## Strengths and limitations of this study

- This systematic review included a wide range of non-operative interventions commonly used in clinical practice.
- This review used consistent inclusion and exclusion criteria for neurogenic claudication, which included the corroboration of a diagnosis of lumbar spinal stenosis with imaging.
- This review used rigorous methods recommended by the Cochrane Back and Neck Pain Review Group including the use of Grading of Recommendations, Assessment, Development and Evaluation to synthesise and summarise the quality of the evidence.
- Only English studies were included in this review.
- Most studies had small samples sizes with heterogeneity in interventions tested, limiting ability to pool data.

**PROSPERO registration number** CRD42020191860.

## INTRODUCTION

Lumbar spinal stenosis (LSS) causing neurogenic claudication is a highly prevalent and rapidly growing public health problem among older adults.<sup>1</sup> It is characterised by bilateral or unilateral buttock pain and/or lower extremity discomfort, pain, weakness or heaviness precipitated by walking and prolonged standing and relieved by stooping forward and sitting.<sup>2 3</sup> The underlying aetiology is usually age-related osteoarthritic changes to lumbar intervertebral discs, facets joints and ligaments leading to narrowing of the central and/or lateral spinal canals and compression and/or ischaemia of the spinal nerves.<sup>2 4</sup>

Limited walking ability is the dominant impairment in neurogenic claudication and the most common reason for seeking care.<sup>5</sup> Limited walking ability due to LSS is associated

with a significant decline in functional status, quality of life and independence in this population.<sup>25</sup>

Although LSS is the most common reason for spine surgery in older adults, most people with neurogenic claudication receive non-operative care.<sup>6</sup> A course of non-operative care is also recommended prior to receiving surgical intervention.<sup>7</sup> However, what constitutes effective non-operative care remains unknown. In 2013, we published a Cochrane review evaluating non-operative treatment for LSS causing neurogenic claudication.<sup>8,9</sup> This review identified 21 randomised controlled trials (RCTs) assessing a variety of non-operative treatments. However, the quality of the evidence was deemed low or very low and therefore no conclusions could be made on the effectiveness of non-operative treatment for neurogenic claudication. The purpose of this study is to update this systematic review and the evidence for non-operative treatments for neurogenic claudication. Our specific research question was: What non-operative interventions are effective in improving outcomes in patients with neurogenic claudication due to LSS?

## METHODS

This systematic review was conducted and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.<sup>10</sup> We used methods recommended by the Cochrane Back Review Group.<sup>11</sup>

### Patient and public involvement statement

Patients or the public were not involved in the conduct of this systematic review.

### Population, interventions, comparison and outcomes criteria

The population of interest was individuals with imaging confirmed LSS (central or foraminal, with or without spondylolisthesis) and neurogenic claudication. Neurogenic claudication is a clinical diagnosis and was defined as buttock or leg pain and/or aching, numbness, tingling, weakness or fatigue with or without back pain, precipitated by standing or walking. There were no age restrictions. The interventions of interest included all non-operative treatments and the comparison was any treatment including surgery. Outcomes included at least one of the following measures: walking ability, pain intensity, physical function, quality of life or global improvement.

### Search and study selection

We replicated and updated our original electronic database search (from 1966 to January 2011) up to 22 July 2020. The search was performed by an experienced librarian in CENTRAL (Cochrane Library 2011 issue1), MEDLINE, EMBASE, CINAHL and Index to Chiropractic Literature databases. The terms 'spinal stenosis', 'lumbar spinal stenosis', 'neurogenic claudication', 'lumbar radicular pain', 'cauda equina' and 'spondylosis' were combined with a highly sensitive search strategy to identify RCTs.

Reference lists of selected studies and previous reviews were also searched to identify additional articles. Online supplemental file 1 provides details on the full search strategies used for all databases.

Studies were included if they were RCTs published in peer-reviewed English journals, at least one arm of the trial provided data on effectiveness of a non-operative treatment and at least 80% of subjects had neurogenic claudication with imaging confirmed LSS. Studies evaluating subjects with radiculopathy caused by disc herniations without neurogenic claudication were excluded.

Studies with mixed populations were only included if separate data for subjects with neurogenic claudication due to LSS were provided.

Two pairs of reviewers independently screened all titles and abstracts identified by the search strategy. Full text of articles deemed to be potentially relevant were independently assessed by two reviewers who made the final decision for inclusion. A third reviewer was consulted if consensus was not reached.

### Risk of bias assessment and data analysis

Two reviewers independently assessed methodological risk of bias and performed data extraction. Safety data (intervention side effects and/or complications) when available were also collected. The Cochrane Risk of Bias Tool 1 was used that included the 12-item criteria recommended by the Cochrane Back Review Group.<sup>11</sup> Discrepancies in risk of bias scoring and data extraction were resolved with discussion and if necessary, with a third reviewer until consensus was reached. Reviewers who were authors of any of the included studies were recused from performing risk of bias assessment, data extraction, data analysis or synthesis of their own studies.

Low risk of bias was defined as fulfilling 6 or more of the 12 criteria including clearly described and appropriate randomisation (item A), and allocation concealment (item B), and with no severe flaws. A severe flaw was defined a priori as a serious methodological deficiency not captured by the 12-item criteria that significantly increases the risk of bias such as very high dropout or cross-over rates and sample sizes less than 30 subjects per treatment arm.

For each comparison, outcomes were analysed according to these follow-up time periods: immediate (up to 1 week following the intervention); short term (between 1 week and 3 months); intermediate (between 3 months and 1 year) and long term (1 year or longer). Outcome data were pooled, and meta-analyses were performed when trials were judged to be sufficiently homogeneous, both clinically and statistically.

Rehabilitation therapy was defined as treatment that used any combination of education, exercise instruction, manual therapy, heat and cold applications, electrotherapy, other physical therapy modalities, orthosis and other assistive devices. Multimodal treatment included various combinations of rehabilitation therapy

treatments, oral and other mediations and spinal injections, but not surgery.

### Data synthesis

The quality of the evidence for each outcome and for each comparison was evaluated using Grading of Recommendations, Assessment, Development and Evaluation (GRADE).<sup>12 13</sup> Overall quality of the evidence was based on performance against five domains: (1) risk of bias; (2) consistency of findings; (3) directness of comparisons; (4) precision of estimates and (5) other considerations such as selective reporting.

The quality of the evidence starts at high when there are consistent findings among at least 75% of RCTs with low risk of bias and consistent, direct and precise data and with no known or suspected publication bias. It downgrades a level for each domain not met. Treatment effects between comparators (more effective, less effective or no difference) were based on statistically significant and clinically important differences in outcomes.

### High-quality evidence

All five domains are met; further research is very unlikely to change the confidence in the estimate of effect.

### Moderate-quality evidence

One of the domains is not met; further research is likely to have an important impact on the confidence in the estimate of effect and may change the estimate.

### Low-quality evidence

Two domains are not met; further research is very likely to have an important impact in the confidence of the estimate of effect and is likely to change the estimate.

### Very low-quality evidence

Three or more domains are not met; there is great uncertainty about the estimate of effect.

Evidence provided by a single small trial was considered inconsistent and imprecise and thus provide 'low' or 'very low' quality evidence, depending on whether it was assessed as having a low or high risk of bias, respectively, and there were no other limitations. Studies with both low risk of bias and inappropriate or unclear randomisation and/or treatment allocation techniques were downgraded by two levels for the 'risk of bias' domain.

The results below are reported based on statistically significant differences between comparators for each outcome using data reported by authors. Differences considered clinically important will be specified when the quality of the evidence is moderate or higher. The minimal clinical important differences (MCIDs) used are listed in the online supplemental table 2. Adverse events for the new studies are detailed when reported by the authors.

## RESULTS

### Selection and description of included trials

We screened 15200 titles and abstracts and assessed 156 full-text articles. This resulted in 44 RCTs meeting the inclusion criteria, including 23 new trials. Figure 1 summarises original and updated screening results. Online supplemental table 1 describes the characteristics of all included trials. In total, 3792 participants (1765 males, 1836 females and 191 participants of undisclosed gender<sup>14 15</sup> were randomised to one of the 60 comparison groups. In total, 17 studies evaluated rehabilitation therapy or multimodal care,<sup>14 16–31</sup> 11 assessed epidural injections,<sup>32–42</sup> 7 evaluated oral medications,<sup>15 43–48</sup> 6 assessed calcitonin,<sup>49–54</sup> 2 evaluated acupuncture<sup>55 56</sup> and 1 assessed spinal manipulation.<sup>57</sup> Thirty-eight trials were conducted at tertiary care or university affiliated centres and six at medical/rehabilitation clinics.<sup>18 24 35–38</sup> The mean age of participants was 63.3 years. The duration of symptoms varied considerably among the studies with a mean ranging from 12 weeks to 15 years. Follow-up periods also varied significantly ranging from immediately following the intervention to 10 years post intervention.

### Risk of bias of included studies

The median and mean number of criteria met was 7 of 12 (range 2–11), see table 1.

Although 31 studies met six or more criteria, only 9 were considered to have low risk of bias.<sup>19 20 24 27 28 31 37 42 43 56</sup> Among the remaining 22 studies that met six or more criteria, 13 failed to explicitly describe and/or use appropriate randomisation procedures, allocation concealment or both<sup>16–18 30 32–34 39 41 48 52 54 57</sup>; 3 had severe flaws due to high cross-over rates,<sup>21 22 25</sup> which made the intention-to-treat analyses uninterpretable and 6 had other serious flaws including premature stopping of the trial,<sup>47</sup> large number of participants lost to follow-up<sup>40</sup> and small sample size (less than 30 participants per arm).<sup>26 29 46 55</sup>


### Evidence of effect of interventions

Overall, 53 of the 60 comparisons were examined in a single trial, most with small sample sizes. It was only possible to combine data from two trials (assessing surgery vs multimodal treatment) for one outcome in a meta-analysis.<sup>19 22</sup> The five other studies (all assessing calcitonin)<sup>49–52 54</sup> were combined qualitatively. The results of these pooled analyses were published in our previous reviews.<sup>8 9</sup> Heterogeneity in source population, intervention and outcome instruments precluded pooling of data from other trials. Online supplemental table 2, summary of GRADE assessment and outcomes, summarises the quality of the evidence for outcomes for each comparison.

### Calcitonin

There were no new studies assessing calcitonin. The conclusion from our previous review was that there is very low-quality evidence from six trials (N=231)<sup>49–54</sup>





**Figure 1** Study flow diagram.

that calcitonin is no better than placebo or paracetamol regardless of mode of administration or outcome assessed.

### Oral medication

We identified four new studies assessing five oral medications. There is low-quality evidence based on one small cross-over trial (N=29)<sup>46</sup> that pregabalin does not improve pain, distance walked, function or global health status immediately following the intervention compared with placebo. Adverse events were reported in 64% of the pregabalin group, the most common being dizziness, compared with 35% in the placebo group.

A small trial evaluating gabapentin plus conservative care (N=45)<sup>48</sup> provides very low-quality evidence demonstrating no significant improvement in back/leg pain, disability scores or global health in the short term compared with conservative care plus botulinum toxin injection. Five patients (20.8%) reported mild-to-moderate pain at injection sites for a few days after botulinum toxin injections.

There is very low-quality evidence from one small trial (N=24)<sup>47</sup> that oxycodone hydrochloride or

propoxyphene and acetaminophen are no better than placebo in the immediate term for all outcomes assessed.

A single small trial provided very low-quality evidence (N=61)<sup>15</sup> that oral corticoids do not improve outcomes in the short term compared with placebo.

The original review identified three studies assessing oral medications and concluded that there is low-quality evidence that prostaglandins improve walking distance and leg pain in the short term compared with etodolac (a non-steroidal anti-inflammatory drug);<sup>43</sup> that there is very low-quality evidence that gabapentin improves walking distance and pain compared with placebo in the intermediate and long term<sup>45</sup> and that methylcobalamin (vitamin B 12) plus conservative treatment improves walking distance in the intermediate and long term compared with conservative treatment alone.<sup>44</sup>

### Rehabilitation therapy and multimodal treatment

We identified eight new studies evaluating 13 rehabilitation therapy and/or multimodal treatment approaches, with one study being compared with surgery.

**Table 1** Risk of bias assessment for studies on non-operative treatment for lumbar spinal stenosis with neurogenic claudication

Author	A	B	C	D	E	F	G	H	I	J	K	L	Total
Calcitonin													
Eskola <i>et al</i> <sup>51</sup>	?	?	+	+	+	?	+	-	?	?	?	+	5
Porter and Hibbert <sup>49</sup>	?	?	-	?	?	+	+	?	-	?	+	+	4
Porter and Miller <sup>50</sup>	?	?	+	?	?	-	+	+	?	?	?	+	4
Podichetty <i>et al</i> <sup>52</sup>	?	?	+	+	+	-	+	-	+	?	?	+	6
Tafazal <i>et al</i> <sup>54</sup>	?	?	+	+	+	+	+	+	-	?	?	+	7
Sahin <i>et al</i> <sup>53</sup>	?	?	-	-	+	-	?	+	+	?	?	+	4
Oral medications													
Prostaglandin													
Matsudaira <sup>43</sup>	+	+	-	-	+	+	?	?	+	?	?	+	7*
Methylcobalamin													
Waikakul and Waikakul <sup>44</sup>	-	?	-	-	+	+	+	?	+	?	?	+	5
Gabapentin													
Yaksi <i>et al</i> <sup>45</sup>	?	?	-	-	-	?	+	+	?	?	?	+	3
Pregabalin													
Markman <i>et al</i> <sup>46</sup>	+	+	+	+	+	+	+	+	?	+	-	+	10†
Gabapentin													
Park <i>et al</i> <sup>48</sup>	+	?	+	+	+	+	+	+	?	?	-	+	8†
Oxymorphone hydrochloride													
Markman <i>et al</i> <sup>47</sup>	+	+	+	+	+	-	?	+	?	+	+	+	9†§
Oral corticoid													
Rodrigues and Natour <sup>15</sup>	+	+	?	?	?	+	+	?	?	?	?	+	5
Rehabilitation therapy or multimodal care													
Goren <i>et al</i> <sup>24</sup>	+	+	-	-	+	+	-	+	+	?	?	+	7*
Koc <i>et al</i> <sup>23</sup>	?	?	-	-	+	+	+	-	+	?	?	+	5
Pua <i>et al</i> <sup>20</sup>	+	+	-	-	+	-	+	+	+	?	-	+	7*
Whitman <i>et al</i> <sup>18</sup>	+	?	-	-	+	+	+	+	+	?	?	+	7
Minelama <i>et al</i> <sup>30</sup>	+	?	-	-	+	+	+	+	?	+	+	+	8†
Schneider <i>et al</i> <sup>31</sup>	+	+	-	-	+	-	+	+	+	?	+	+	8*
Ammendolia <i>et al</i> <sup>27</sup>	+	+	-	-	+	+	+	+	+	+	+	+	10*
Oğuz <i>et al</i> <sup>14</sup>	?	?	-	-	?	?	+	-	?	?	?	+	2
Homayouni <i>et al</i> <sup>26</sup>	+	+	-	-	+	+	+	-	-	?	?	+	7†
Marchand <i>et al</i> <sup>29</sup>	+	+	-	-	+	?	+	+	?	-	+	+	7†
Kim <i>et al</i> <sup>28</sup>	+	+	+	+	+	+	+	+	?	+	+	+	11*
Spinal manipulation													
Passmore <i>et al</i> <sup>67</sup>	-	+	-	-	+	+	+	-	+	+	+	+	8†

Continued

Table 1 Continued

Author	A	B	C	D	E	F	G	H	I	J	K	L	Total
Acupuncture													
Kim <i>et al</i> <sup>55</sup>	+	+	-	-	-	-	+	+	-	+	+	+	7†
Qin <i>et al</i> <sup>56</sup>	+	+	+	-	+	+	+	+	+	-	+	+	10*
Epidural injections													
Cuckler <i>et al</i> <sup>32</sup>	?	?	+	+	+	+	+	+	+	?	+	+	9
Fukusaki <i>et al</i> <sup>34</sup>	?	?	?	?	+	+	+	+	+	?	+	+	7
Zahaar <sup>33</sup>	?	?	+	?	+	+	+	+	+	-	?	-	6
Brown <sup>35</sup>	+	-	+	-	?	+	+	-	?	?	-	+	5
Friedly <i>et al</i> <sup>37</sup> , Friedly <i>et al</i> <sup>62</sup> and Makris <sup>64</sup>	+	+	+	+	+	+	+	+	?	+	+	+	11*
Song <i>et al</i> <sup>36</sup>	?	?	?	?	?	+	+	-	?	+	+	+	5
Milburn <i>et al</i> <sup>38</sup>	?	?	+	-	+	-	+	-	?	-	-	+	4
Hammerich <i>et al</i> <sup>40</sup>	+	+	-	-	+	-	+	?	?	-	+	+	6†
Sencan <i>et al</i> <sup>41</sup>	+	?	+	-	+	+	?	+	+	+	?	+	8†
Wei <i>et al</i> <sup>42</sup>	+	+	+	-	-	+	-	+	?	+	+	+	8*
Percutaneous epidural adhesiolysis													
Karm <i>et al</i> <sup>39</sup>	+	?	+	-	+	-	+	+	?	-	-	+	6†
Surgery versus physical therapy													
Zucherman <i>et al</i> <sup>17</sup>	?	+	-	-	+	+	+	+	+	?	+	+	>6
Zucherman <i>et al</i> <sup>60</sup> and Hsu <i>et al</i> <sup>69</sup>													
Weinstein <i>et al</i> <sup>21</sup> , Weinstein <i>et al</i> <sup>68</sup> , Abdu <i>et al</i> <sup>61</sup>	+	+	-	-	+	+	+	+	?	?	-	+	>9***††
Amundsen <i>et al</i> <sup>16</sup>	+	?	-	-	-	+	+	+	-	?	-	?	4
Malmivaara <i>et al</i> <sup>19</sup>	+	+	-	-	+	+	+	+	+	?	?	+	8*
Weinstein <i>et al</i> <sup>22</sup> , Weinstein <i>et al</i> <sup>69</sup> , Lurie <i>et al</i> <sup>66</sup>	+	+	-	-	+	-	+	+	?	?	-	+	6††
Delitto <i>et al</i> <sup>25</sup>	+	+	-	-	+	?	+	-	+	-	+	+	7††

A: Was the method of randomisation adequate? B: Was the treatment allocation concealed? C: Was the patient blinded to the intervention? D: Was the care provider blinded to the intervention? E: Was the outcome assessor blinded to the intervention? F: Was the drop-out rate described and acceptable? G: Were all randomised participants analysed in the group to which they were allocated? H: Are reports of the study free of suggestion of selective outcome reporting? I: Were the groups similar at baseline regarding the most important prognostic indicators? J: Were co-interventions avoided or similar? K: Was the compliance acceptable in all groups? L: Was the timing of the outcome assessment similar in all groups? +: Yes, -: No, ?: Unclear.

\*Low risk of bias if six or more items met, including valid randomisation and treatment allocation techniques and no severe flaws.

†<30 participants per treatment arm.

‡Treatment allocation unclear.

\$Premature end of study.

||2-year follow-up drop-out rate 30%, 1 year<20%; intention to treat inconsistent at 2-year follow-up.

\*\*Drop-out rate<20% at 1 year, >20% at 4 years.

††Severe flaw due to high cross-over rates.

There is moderate-quality evidence from one trial (N=259)<sup>31</sup> that manual therapy and exercise provides superior and clinically important short-term improvement in symptoms and function compared with medical care or community-based group exercise and that community-based group exercise improves physical activity in the short term compared with medical care. There were no reported serious adverse events in any group. There was a significantly greater rate of transient joint soreness associated with the manual therapy and exercise group (49%) compared with the community-based group exercise (31%) and medical care (6%) groups.

Another trial provides moderate-quality evidence (N=104)<sup>27</sup> that comprehensive care (manual therapy, education and exercise delivered using a cognitive-behavioural approach) demonstrates superior and clinically important improvements in walking distance in the immediate, short, intermediate and long term and compared with self-directed home exercise. This study also provides low-quality evidence that comprehensive care improves overall pain and function in the long term compared with self-directed home exercises. At 12 months, none of the 43 participants in the comprehensive group and 2 of the 46 participants in the self-directed group experienced adverse events. These adverse events were mostly attributed to a temporary increase in low back and/or leg pain.

There is low-quality evidence from one trial (N=34)<sup>28</sup> that a form of manual therapy (Mokuri Chuna), acupuncture and physician care, with or without a herbal remedy (Gang-Chuk Tang), improves low back pain in the intermediate term compared with oral aceclofenac, epidural steroids and physical therapy (heat and TENS).

A single study assessing supervised physical therapy (manual therapy, exercise and body weight-supported treadmill) (N=86)<sup>30</sup> provides low-quality evidence for improved symptoms, function and walking distance in the short term compared with home exercises.

There is very low-quality evidence from one study (N=120)<sup>14</sup> that heat, transcutaneous electrical nerve stimulation (TENS) and home exercise instruction are no better than isokinetic exercise in the immediate, short and intermediate term for all outcomes and less effective than unloaded exercises in the immediate and short term. Unloaded exercise was also found to be superior to isokinetic exercise in the immediate and short term.

One small single study (N=47)<sup>26</sup> provides very low-quality evidence that aquatic exercise is more effective than physical therapy (exercise, ultrasound, heat and TENS) in improving pain and walking distance in the immediate term.

Another small single trial (N=40)<sup>29</sup> provides very low-quality evidence that a presurgical exercise programme improves postsurgical outcomes in the immediate, but not in the short or intermediate terms.

There is low-quality evidence from one study (N=169)<sup>25</sup> that a structured physical therapy programme (education

and exercises) provides similar outcomes to decompression surgery in the long term (2-year follow-up). Overall, 9 out of 82 participants receiving physical therapy reported adverse events consisting of worsening of symptoms, whereas 33 out of 87 participants reported surgery-related complications, mainly attributable to reoperation, delay in wound healing and surgical site infection.

Our original review identified nine rehabilitation therapy/multimodal trials of which five were compared with surgical interventions. A meta-analysis was conducted for two of the surgical trials. Two of the original surgical trials have since published 8-year follow-up results (see below). All studies provide either low-quality or very low-quality evidence.

A meta-analysis<sup>8 9</sup> that includes two trials<sup>22 19</sup> shows that laminectomy improves outcomes only at the 2-year follow-up compared with conservative care. One of these studies shows no difference in outcomes after an 8-year follow-up.<sup>58</sup>

An interspinous surgical implant<sup>17 59 60</sup> was found to be superior to multimodal treatment (epidural injections, pain medication, education, exercise, back brace, heat/ice and massage). Another trial<sup>16</sup> provided inconclusive evidence when comparing laminectomy with or without fusion to lumbar orthosis and education.

Among patients with degenerative spondylolisthesis, one study<sup>21</sup> shows no difference in outcomes with laminectomy when compared with conservative care, including after an 8-year follow-up.<sup>61</sup>

One study showed that exercise plus ultrasound is no better than exercise plus sham ultrasound but better than no treatment, and exercise plus sham ultrasound is better than no treatment.<sup>24</sup> Other studies demonstrated that inpatient physical therapy (ultrasound, heat and TENS) is more effective than home exercise plus oral diclofenac,<sup>23</sup> unweighted treadmill walking plus exercise is no better than cycling plus exercise<sup>20</sup> and manual therapy, exercise and unweighted treadmill are more effective than flexion exercises, walking and sham ultrasound.<sup>18</sup>

### Epidural injections

We identified six new studies evaluating epidural injections. There is moderate-quality evidence from one study (N=400)<sup>37 62</sup> that glucocorticoid plus lidocaine injection is better than lidocaine alone in improving pain and function at 3 weeks (short term) but not at 6 weeks (short term), 12 weeks (intermediate term) or 12 months (long term). The improved outcomes at 3 weeks were statistically significant but not considered to be of clinical importance.<sup>63</sup> A follow-up subgroup analysis<sup>64</sup> using patient-prioritised Roland-Morris Disability Questionnaire items did not change the results. A total 21.5% of patients in the glucocorticoid-lidocaine group and 15.5% in the lidocaine alone group reported one or more adverse events (p=0.08). Adverse events included headaches, fever, infection, dizziness, cardiovascular/lung problems, leg swelling and dural puncture.

A small study (N=29)<sup>36</sup> provided very low-quality evidence that an injection of lidocaine is no better than a saline injection for all outcomes in the short term.

There is very low-quality evidence from one study (N=57)<sup>38</sup> that steroid injections at the level of maximal stenosis improve pain and function in the immediate and short term compared with steroid injections at two levels cephalad to the maximum level of stenosis.

A small trial (N=54)<sup>40</sup> provided very low-quality evidence that steroid injections are no better than steroid injections combined with physical therapy (manual therapy and exercise) in improving pain or function in the short term but are more effective in improving pain in the intermediate and long term.

There is very low-quality evidence from one study (N=67)<sup>41</sup> that interlaminar steroid injection improves pain and walking distance in the intermediate but not in the short term compared with transforaminal steroid injection.

A three-arm trial (N=30)<sup>42</sup> provided low-quality evidence that tumour necrosis factor (TNF) alpha inhibitor (etanercept) injections improved pain and function in the immediate, short and intermediate term compared with steroid or lidocaine injections and that steroid injections were no better than lidocaine for all outcomes and follow-up periods.

There is very low-quality evidence from one small trial (N=38)<sup>35</sup> that minimally invasive lumbar decompression surgery is no better than epidural steroid injections for all outcomes in the short term.

One small trial (N=44)<sup>39</sup> provided very low-quality evidence that an epidural inflatable balloon catheter (ZiNeu) improves pain and function in the intermediate term but not in the short term compared with a balloon-less catheter (Racz). Minor and transient adverse events were reported equally in both groups (no data provided), mostly pain and paraesthesia at the injection site.

Our original review identified four trials evaluating seven epidural injection approaches, all with very low-quality evidence for all outcomes. Two trials demonstrated that translaminar<sup>32</sup> or caudal<sup>33</sup> steroid injections were no better than placebo. Two other trials showed that translaminar epidural steroid plus a block was better than placebo or an epidural block alone,<sup>34</sup> that translaminar epidural block was better than placebo<sup>34</sup> and that interlaminar epidural steroid plus a block was better than home exercise plus diclofenac or inpatient physical therapy (ultrasound, heat and TENS).<sup>23</sup>

### Acupuncture

We identified two new studies assessing acupuncture. There is low-quality evidence from one trial (N=80)<sup>56</sup> that acupuncture improves back and leg pain, symptoms and function in the immediate, short and intermediate term compared with sham acupuncture. Overall, 3 out of 40 participants in the acupuncture group reported short-term pain at the insertion site (one also had a haematoma) and 5 out of the 40 participants in the sham group

reported non-serious back pain or fatigue. There is very low-quality evidence from a small trial (N=50)<sup>55</sup> that acupuncture plus usual care is no better than usual care alone in the short term for all outcomes.

### Spinal manipulation

We identified one study assessing spinal manipulation. There is very low-quality evidence from a very small trial (N=14)<sup>57</sup> that spinal manipulation alone is no better than a wait list control in the immediate term for all outcomes.

## DISCUSSION

We updated our systematic review on non-operative treatments for LSS causing neurogenic claudication and identified 23 new trials that were added to the previous 21 studies. The highest number of studies, 17/44, evaluated rehabilitation therapy/multimodal treatment, 11 assessed epidural interventions, 7 assessed oral medications, 6 assessed calcitonin, 2 evaluated acupuncture and 1 assessed spinal manipulation. Of the 60 comparisons that were evaluated, 5 comparisons from three trials<sup>27 31 37</sup> provided moderate-quality evidence. The remaining comparisons provide either low-quality or very low-quality evidence. In our original review, all comparisons for all the interventions assessed were of low-quality or very low-quality evidence. This lack of moderate-quality or high-quality evidence limited our ability to make conclusions on the effectiveness of most non-operative treatments.

There is now moderate evidence that a multimodal structured 6-week programme consisting of manual therapy and exercise with or without education is an effective treatment approach<sup>27 31</sup> for neurogenic claudication and that epidural steroid injections do not provide clinically important improvements in short-term or long-term outcomes compared with epidural lidocaine injections. However, given that these respective findings came from single studies, this evidence lacks consistency and therefore there is a possibility that replicating these trials in the future might result in substantially different conclusions. However, a recent clinical practice guideline for the management of LSS leading to neurogenic claudication concurred with our findings and recommended, based on moderate-quality evidence, multimodal care consisting of education with home exercises and manual therapy.<sup>65</sup> These guidelines also recommended against the use of epidural steroid injections, based on high-quality evidence. A recent systematic review and meta-analysis of RCTs evaluating conservative non-pharmacological therapies for degenerative LSS also concluded, based on low-to-moderate evidence, that manual therapy and supervised exercises significantly improve outcomes compared with self-directed or group exercises.<sup>66</sup> A recent clinical update published in the *British Medical Journal* recommended supervised exercise and manual therapy as a first-line treatment for LSS and recommended against the use of epidural steroid injections.<sup>67</sup> More dated systematic



reviews did not recommend a combination of education, exercise and manual therapy as an effective treatment for LSS.<sup>7 68 69</sup> However, these reviews did not include the more recent higher quality trials<sup>27 31</sup> evaluating this multimodal approach.

A multimodal approach to the treatment of LSS would appear to be a rational approach given the complexity of neurogenic claudication with underlying physical, functional and psychosocial factors impacting recovery.<sup>70</sup> There is also a plausible rationale for the lack of effectiveness of epidural steroid injections for neurogenic claudication since the dominant underlying pathophysiological mechanism appears to be neuroischaemia rather than neuroinflammation.<sup>4</sup>

Although we cannot make firm conclusions about the effectiveness of non-operative treatments for neurogenic claudication, this review is important because it provides important information regarding the state of current evidence regarding non-operative treatments. This can be used to inform clinical practice guidelines and aid clinicians and patients in making clinical decisions regarding treatment options. This is particularly important with respect to interventions that have higher risks and costs such as epidural injections and surgery. About 25% of all epidural injections are performed for LSS<sup>71 72</sup> yet the evidence from our current review and those of others<sup>73–75</sup> do not support their use. The number and associated costs of surgical procedures for degenerative LSS are growing, especially decompression surgery with complex fusion.<sup>76 77</sup> LSS continues to be the most common reason for spine surgery in older adults.<sup>6 76</sup> High-quality evidence for the effectiveness of surgery is also lacking based on our current review and the findings of other systematic reviews.<sup>78 79</sup> Clinical trials evaluating surgery for LSS are difficult to conduct due to challenges in recruitment and blinding (patient and practitioner) and high costs.<sup>80</sup> One ongoing clinical trial is comparing decompression surgery with sham surgery which should help to evaluate the potential role of the placebo effect of surgery for LSS.<sup>81</sup>

Oral medication is often the first-line treatment in primary care management of LSS.<sup>5</sup> Pregabalin and gabapentin are commonly prescribed medications for LSS despite the growing evidence that these medications are not effective for back-related leg symptoms and may cause more harm than good.<sup>82–84</sup>

New to this updated review are clinical trials on acupuncture and spinal manipulation; however, the quality of the evidence was insufficient to make conclusions on their effectiveness. A systematic review and meta-analysis of RCTs and controlled clinical trials published in Chinese found no conclusive evidence for the effectiveness and safety of acupuncture for LSS.<sup>85</sup> Passive unimodal treatments such as acupuncture and spinal manipulation are unlikely to provide long-term benefit but more likely to provide benefit when combined with a comprehensive approach to managing LSS,<sup>27</sup> not unlike recommendations for managing chronic low back pain.<sup>86</sup>

This review is also important because it provides a comprehensive assessment and identification of significant knowledge gaps in this area to guide future research. This includes the need for higher quality studies that assess commonly used non-operative treatments particularly in primary care settings that are adequately powered and have low risk of bias and long-term follow-up. Future RCTs should follow the Consolidated Standards of Reporting Trials guideline<sup>87</sup> when planning trials and reporting study findings in an attempt to improve transparency and reduce bias.

The strengths of this review include the evaluation of a wide range of non-operative interventions and the use of consistent inclusion and exclusion criteria for neurogenic claudication, which included the corroboration of a diagnosis of LSS with imaging. The use of these criteria to define the study population increases the likelihood that participants in the included studies had the diagnosis of neurogenic claudication due to narrowing of the central canal or lateral foraminae.<sup>88–90</sup> Other strengths of this review include the use of rigorous methods recommended by the Cochrane Collaboration, the WHO and the Cochrane Back and Neck Pain Review Group.<sup>13</sup> This included the use of the GRADE method to synthesise and summarise the quality of the evidence.

Limitations of this review include the potential for language bias because only English articles were accepted. We also included studies with small samples sizes which are more prone to high risk of bias.<sup>91</sup> Over half of the included studies had less than 30 subjects per arm at baseline and none of these studies could be pooled because of high heterogeneity across studies. However, the exclusion of studies with small samples sizes in this review would not have changed our conclusions. The definition of a severe flaw and the cut-off point of 6 or more to differentiate trials of low from high risk of bias were arbitrary, therefore alternative definitions and cut-off points or the use of other risk of bias tools could have impacted the findings and conclusions of this review. The validity of MCIDs used in this review is unknown. Although most were derived from studies with neurogenic claudication,<sup>63 92 93</sup> others were based on an arbitrary improvement of at least 30%.<sup>94</sup> There are no agreed upon MCIDs in LSS and therefore different MCIDs thresholds could have potentially altered our conclusions. The location and severity of the stenosis on imaging was not deemed important in this review. Imaging findings often do not correlate with patient symptoms or severity and therefore imaging by itself is a not reliable diagnostic tool in this population.<sup>67 95 96</sup> Neurogenic claudication is the clinical entity of interest in this review and, although usually caused by LSS, the diagnosis is made clinically without imaging.<sup>97</sup> Neurogenic claudication symptoms, by definitions, improve with flexion, due to the increased volume around the involved nerve roots irrespective of where the stenosis is located (eg, centrally or at the lateral recess). However, it is uncertain whether the effectiveness of some interventions, such as epidural steroid injections, is dependent on

location of the spinal stenosis. This is a different research question requiring future research.

## CONCLUSIONS

There is moderate-quality evidence that a multimodal approach that includes manual therapy and exercise, with or without education, is a safe and effective treatment and that epidural steroids are not effective for the management of LSS causing neurogenic claudication. All other studies evaluating non-operative interventions provided insufficient quality evidence, limiting the ability to make conclusions about their effectiveness. With the growing prevalence and significant personal, social and economic burden of LSS, more high-quality evidence for non-operative interventions is urgently needed to guide clinical practice.

## Author affiliations

<sup>1</sup>Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada

<sup>2</sup>Rheumatology, Sinai Health System, Toronto, Ontario, Canada

<sup>3</sup>Graduate Education and Research, Canadian Memorial Chiropractic College, Toronto, Ontario, Canada

<sup>4</sup>School of Physical and Occupational Therapy, Faculty of Medicine and Health Sciences, McGill University, Montreal, Quebec, Canada

<sup>5</sup>Département Chiropratique, Université du Québec à Trois-Rivières, boulevard des Forges, Trois-Rivières Québec, Canada

<sup>6</sup>Physical Therapy, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

<sup>7</sup>Sports Medicine and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark

<sup>8</sup>Toronto Rehabilitation Institute, Toronto, Ontario, Canada

<sup>9</sup>Institute for Work & Health, Toronto, Ontario, Canada

<sup>10</sup>Faculty of Health Sciences, Ontario Tech University, Oshawa, Ontario, Canada

<sup>11</sup>Health Systems Management, Rush University, Chicago, Illinois, USA

**Twitter** Andrea D Furlan @adfurlan

**Acknowledgements** The authors would like to thank librarian Maggie Tiong from the Institute for Work & Health for her assistance in developing the search strategy and retrieving relevant articles and Carly Coelho for assistance in screening titles and abstracts.

**Contributors** CA acted as the study guarantor and was involved in the conception and design of the study, screening of articles, risk of bias assessment, Grading of Recommendations, Assessment, Development and Evaluation (GRADE) analysis, writing the first draft of the manuscript, revision of the manuscript and administrative support. AB, MJS, ADF, CC and JO were involved in screening of articles, risk of bias assessment, GRADE analysis and critical revision of the manuscript. CH, JP, AA, KS, JY and AAdeboyejo participated in screening of articles, risk of bias assessment, data extraction and critical revision of the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** CA received research funding from the Canadian Chiropractic Research Foundation and the Arthritis Society. JY has received funding support from the Danish Foundation for Chiropractic Research and Postgraduate Education, the Ontario Chiropractic Association, the Canadian Memorial Chiropractic College, the National Chiropractic Mutual Insurance Company Foundation and the University of Southern Denmark. CC holds a Research Chair in Knowledge Translation in the Faculty of Health Sciences, Ontario Tech University, funded by the Canadian Chiropractic Research Foundation.

**Patient consent for publication** Not applicable.

**Ethics approval** This study does not involve human participants.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplementary information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Carlo Ammendolia <http://orcid.org/0000-0003-4417-4961>

André Bussi eres <http://orcid.org/0000-0002-2818-6949>

James J Young <http://orcid.org/0000-0003-1210-3106>

Andrea D Furlan <http://orcid.org/0000-0001-6138-8510>

## REFERENCES

- Jensen RK, Jensen TS, Koes B, *et al.* Prevalence of lumbar spinal stenosis in general and clinical populations: a systematic review and meta-analysis. *Eur Spine J* 2020;29:2143–63.
- Katz JN, Harris MB, *et al.* Lumbar spinal stenosis. *N Engl J Med* 2008;358:818–25.
- Comer CM, Redmond AC, Bird HA, *et al.* Assessment and management of neurogenic claudication associated with lumbar spinal stenosis in a UK primary care musculoskeletal service: a survey of current practice among physiotherapists. *BMC Musculoskelet Disord* 2009;10:121.
- Kobayashi S, Pathophysiology KS. Pathophysiology, diagnosis and treatment of intermittent claudication in patients with lumbar canal stenosis. *World J Orthop* 2014;5:134–45.
- Fritz JM, Delitto A, Welch WC, *et al.* Lumbar spinal stenosis: a review of current concepts in evaluation, management, and outcome measurements. *Arch Phys Med Rehabil* 1998;79:700–8.
- Deyo RA. Treatment of lumbar spinal stenosis: a balancing act. *Spine J* 2010;10:625–7.
- Rousing R, Jensen RK, Fruensgaard S, *et al.* Danish national clinical guidelines for surgical and nonsurgical treatment of patients with lumbar spinal stenosis. *Eur Spine J* 2019;28:1386–96.
- Ammendolia C, Stuber K, de Bruin LK, *et al.* Nonoperative treatment of lumbar spinal stenosis with neurogenic claudication: a systematic review. *Spine* 2012;37:E609–16.
- Ammendolia C, Stuber KJ, Rok E, *et al.* Nonoperative treatment for lumbar spinal stenosis with neurogenic claudication. *Cochrane Database Syst Rev* 2013;8:CD010712.
- Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010;8:336–41.
- Furlan AD, Pennick V, Bombardier C, *et al.* 2009 updated method guidelines for systematic reviews in the Cochrane back review group. *Spine* 2009;34:1929–41.
- Atkins D, Eccles M, Flottorp S, *et al.* Systems for grading the quality of evidence and the strength of recommendations I: critical appraisal of existing approaches the grade Working group. *BMC Health Serv Res* 2004;4:38.
- Furlan AD, Malmivaara A, Chou R, *et al.* 2015 updated method guideline for systematic reviews in the Cochrane back and neck group. *Spine* 2015;40:1660–73.
- O uz H, Levendo lu F, Karahan AY. Comparison of effects of standard, isokinetic and unloading exercises in the treatment of lumbar spinal stenosis. *Turk J Phys Med Rehab* 2013;16:1–7.
- Rodrigues LCL, Natour J, A double-blind NJ. A double-blind, randomized controlled, prospective trial assessing the effectiveness of oral corticoids in the treatment of symptomatic lumbar canal stenosis. *J Negat Results Biomed* 2014;13:13:13.
- Amundsen T, Weber H, Nordal HJ, *et al.* Lumbar spinal stenosis: conservative or surgical management?: a prospective 10-year study. *Spine* 2000;25:1424–35. discussion 35–6.
- Zucherman JF, Hsu KY, Hartjen CA, *et al.* A prospective randomized multi-center study for the treatment of lumbar spinal stenosis

- with the X stop interspinous implant: 1-year results. *Eur Spine J* 2004;13:22–31.
- 18 Whitman JM, Flynn TW, Childs JD, *et al.* A comparison between two physical therapy treatment programs for patients with lumbar spinal stenosis: a randomized clinical trial. *Spine* 2006;31:2541–9.
  - 19 Malmivaara A, Slätis P, Heliövaara M, *et al.* Surgical or nonoperative treatment for lumbar spinal stenosis? A randomized controlled trial. *Spine* 2007;32:1–8.
  - 20 Pua Y-H, Cai C-C, Lim K-C. Treadmill walking with body weight support is no more effective than cycling when added to an exercise program for lumbar spinal stenosis: a randomised controlled trial. *Aust J Physiother* 2007;53:83–9.
  - 21 Weinstein JN, Lurie JD, Tosteson TD, *et al.* Surgical versus nonsurgical treatment for lumbar degenerative spondylolisthesis. *N Engl J Med* 2007;356:2257–70.
  - 22 Weinstein JN, Tosteson TD, Lurie JD, *et al.* Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med* 2008;358:794–810.
  - 23 Koc Z, Ozcakar S, Sivrioğlu K, *et al.* Effectiveness of physical therapy and epidural steroid injections in lumbar spinal stenosis. *Spine* 2009;34:985–9.
  - 24 Goren A, Yildiz N, Topuz O, *et al.* Efficacy of exercise and ultrasound in patients with lumbar spinal stenosis: a prospective randomized controlled trial. *Clin Rehabil* 2010;24:623–31.
  - 25 Delitto A, Piva SR, Moore CG, *et al.* Surgery versus nonsurgical treatment of lumbar spinal stenosis: a randomized trial. *Ann Intern Med* 2015;162:465–73.
  - 26 Homayouni K, Naseri M, Zaravar F, *et al.* Comparison of the effect of aquatic physical therapy and conventional physical therapy in patients with lumbar spinal stenosis (a randomized controlled trial). *J Musculoskelet Res* 2015;18:1550002.
  - 27 Ammendolia C, Côté P, Southerst D, *et al.* Comprehensive nonsurgical treatment versus Self-directed care to improve walking ability in lumbar spinal stenosis: a randomized trial. *Arch Phys Med Rehabil* 2018;99:2408–19.
  - 28 Kim K, Shin K-M, Hunt CL, *et al.* Nonsurgical integrative inpatient treatments for symptomatic lumbar spinal stenosis: a multi-arm randomized controlled pilot trial. *J Pain Res* 2019;12:1103–13.
  - 29 Marchand A-A, Sutin M, O'Shaughnessy J, *et al.* Feasibility of conducting an active exercise prehabilitation program in patients awaiting spinal stenosis surgery: a randomized pilot study. *Sci Rep* 2019;9:12257.
  - 30 Minetama M, Kawakami M, Teraguchi M, *et al.* Supervised physical therapy vs. home exercise for patients with lumbar spinal stenosis: a randomized controlled trial. *Spine J* 2019;19:1310–8.
  - 31 Schneider MJ, Ammendolia C, Murphy DR, *et al.* Comparative clinical effectiveness of nonsurgical treatment methods in patients with lumbar spinal stenosis: a randomized clinical trial. *JAMA Netw Open* 2019;2:e186828.
  - 32 Cuckler JM, Bernini PA, Wiesel SW, *et al.* The use of epidural steroids in the treatment of lumbar radicular pain. A prospective, randomized, double-blind study. *J Bone Joint Surg Am* 1985;67:63–6.
  - 33 Zahaar MS. The value of caudal epidural steroids in the treatment of lumbar neural compression syndromes. *J Neurol Orthop Med Surg* 1991;12:181–4.
  - 34 Fukusaki M, Kobayashi I, Hara T, *et al.* Symptoms of spinal stenosis do not improve after epidural steroid injection. *Clin J Pain* 1998;14:148–51.
  - 35 Brown LL, double-blind A. A double-blind, randomized, prospective study of epidural steroid injection vs. the mild® procedure in patients with symptomatic lumbar spinal stenosis. *Pain Pract* 2012;12:333–41.
  - 36 Song SH, Ryu GH, Park JW, *et al.* The effect and safety of steroid injection in lumbar spinal stenosis: with or without local anesthetics. *Ann Rehabil Med* 2016;40:14–20.
  - 37 Friedly JL, Comstock BA, Turner JA, *et al.* A randomized trial of epidural glucocorticoid injections for spinal stenosis. *N Engl J Med* 2014;371:11–21.
  - 38 Milburn J, Freeman J, Steven A, *et al.* Interlaminar epidural steroid injection for degenerative lumbar spinal canal stenosis: does the intervertebral level of performance matter? *Ochsner J* 2014;14:62–6.
  - 39 Karm M-H, Choi S-S, Kim D-H, *et al.* Percutaneous epidural adhesiolysis using inflatable balloon catheter and balloonless catheter in central lumbar spinal stenosis with neurogenic claudication: a randomized controlled trial. *Pain Physician* 2018;21:593–606.
  - 40 Hammerich A, Whitman J, Mintken P, *et al.* Effectiveness of physical therapy combined with epidural steroid injection for individuals with lumbar spinal stenosis: a randomized parallel-group trial. *Arch Phys Med Rehabil* 2019;100:797–810.
  - 41 Sencan S, Edipoglu IS, Celenioglu AE, *et al.* Comparison of treatment outcomes in lumbar central stenosis patients treated with epidural steroid injections: interlaminar versus bilateral transforaminal approach. *Korean J Pain* 2020;33:226–33.
  - 42 Wei P, Xu Y, Yao Q, *et al.* Randomized trial of 3-drug combination for lumbar nerve root epidural injections with a TNF- $\alpha$  inhibitor in treatment of lumbar stenosis. *Br J Neurosurg* 2020;34:168–71.
  - 43 Matsudaira K, Seichi A, Kunogi J, *et al.* The efficacy of prostaglandin E1 derivative in patients with lumbar spinal stenosis. *Spine* 2009;34:115–20.
  - 44 Waikakul W, Waikakul S. Methylcobalamin as an adjuvant medication in conservative treatment of lumbar spinal stenosis. *J Med Assoc Thai* 2000;83:825–31.
  - 45 Yaksi A, Özgönenel L, Özgönenel B. The efficiency of gabapentin therapy in patients with lumbar spinal stenosis. *Spine* 2007;32:939–42.
  - 46 Markman JD, Frazer ME, Rast SA, *et al.* Double-Blind, randomized, controlled, crossover trial of pregabalin for neurogenic claudication. *Neurology* 2015;84:265–72.
  - 47 Markman JD, Gewandter JS, Frazer ME, *et al.* A randomized, double-blind, placebo-controlled crossover trial of oxymorphone hydrochloride and propoxyphene/acetaminophen combination for the treatment of neurogenic claudication associated with lumbar spinal stenosis. *Spine* 2015;40:684–91.
  - 48 Park SJ, Yoon KB, Yoon DM, *et al.* Botulinum toxin treatment for nocturnal calf cramps in patients with lumbar spinal stenosis: a randomized clinical trial. *Arch Phys Med Rehabil* 2017;98:957–63.
  - 49 Porter RW, Hibbert C. Calcitonin treatment for neurogenic claudication. *Spine* 1983;8:585–92.
  - 50 Porter RW, Miller CG. Neurogenic claudication and root claudication treated with calcitonin. A double-blind trial. *Spine* 1988;13:1061–4.
  - 51 Eskola A, Pohjolainen T, Alaranta H, *et al.* Calcitonin treatment in lumbar spinal stenosis: a randomized, placebo-controlled, double-blind, cross-over study with one-year follow-up. *Calcif Tissue Int* 1992;50:400–3.
  - 52 Podichetty VK, Segal AM, Lieber M, *et al.* Effectiveness of salmon calcitonin nasal spray in the treatment of lumbar canal stenosis: a double-blind, randomized, placebo-controlled, parallel group trial. *Spine* 2004;29:2343–9.
  - 53 Sahin F, Yilmaz F, Kotevoglou N, *et al.* The efficacy of physical therapy and physical therapy plus calcitonin in the treatment of lumbar spinal stenosis. *Yonsei Med J* 2009;50:683–8.
  - 54 Tafazal SI, Ng L, Sell P. Randomised placebo-controlled trial on the effectiveness of nasal salmon calcitonin in the treatment of lumbar spinal stenosis. *Eur Spine J* 2007;16:207–12.
  - 55 Kim KH, Kim YR, Baik SK, *et al.* Acupuncture for patients with lumbar spinal stenosis: a randomised pilot trial. *Acupunct Med* 2016;34:267–74.
  - 56 Qin Z, Ding Y, Xu C, *et al.* Acupuncture vs noninsertive sham acupuncture in aging patients with degenerative lumbar spinal stenosis: a randomized controlled trial. *Am J Med* 2020;133:e20:500–7.
  - 57 Passmore SR, Johnson MG, Aloraini SM, *et al.* Impact of spinal manipulation on lower extremity motor control in lumbar spinal stenosis patients: a small-scale assessor-blind randomized clinical trial. *J Manipulative Physiol Ther* 2019;42:23–33.
  - 58 Lurie JD, Tosteson TD, Tosteson A, *et al.* Long-Term outcomes of lumbar spinal stenosis: eight-year results of the spine patient outcomes research trial (sport). *Spine* 2015;40:63–76.
  - 59 Hsu KY, Zucherman JF, Hartjen CA, *et al.* Quality of life of lumbar stenosis-treated patients in whom the X stop interspinous device was implanted. *J Neurosurg Spine* 2006;5:500–7.
  - 60 Zucherman JF, Hsu KY, Hartjen CA, *et al.* A multicenter, prospective, randomized trial evaluating the X stop interspinous process decompression system for the treatment of neurogenic intermittent claudication: two-year follow-up results. *Spine* 2005;30:1351–8.
  - 61 Abdu WA, Sacks OA, Tosteson ANA, *et al.* Long-term results of surgery compared with nonoperative treatment for lumbar degenerative spondylolisthesis in the spine patient outcomes research trial (sport). *Spine* 2018;43:1619–30.
  - 62 Friedly JL, Comstock BA, Turner JA, *et al.* Long-Term effects of repeated injections of local anesthetic with or without corticosteroid for lumbar spinal stenosis: a randomized trial. *Arch Phys Med Rehabil* 2017;98:1499–507.
  - 63 Cleland JA, Whitman JM, Houser JL, *et al.* Psychometric properties of selected tests in patients with lumbar spinal stenosis. *Spine J* 2012;22:921–31.
  - 64 Makris UE, Edwards TC, Lavalley DC, *et al.* Patient priority weighting of the Roland Morris disability questionnaire does not change results of the lumbar epidural steroid injections for spinal stenosis trial. *Spine* 2017;42:42–8.



- 65 Bussi res A, Cancelliere C, Ammendolia C, *et al.* Non-Surgical interventions for lumbar spinal stenosis leading to neurogenic claudication: a clinical practice guideline. *J Pain* 2021;22:1015–39.
- 66 Jacobi S, Beynon A, Dombrowski SU, *et al.* Effectiveness of conservative nonpharmacologic therapies for pain, disability, physical capacity, and physical activity behavior in patients with degenerative lumbar spinal stenosis: a systematic review and meta-analysis. *Arch Phys Med Rehabil* 2021;102:2247–60.
- 67 Jensen RK, Harhangi BS, Huygen F, *et al.* Lumbar spinal stenosis. *BMJ* 2021;61:n1581.
- 68 Kreiner DS, Shaffer WO, Baisden JL, *et al.* An evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spinal stenosis (update). *Spine J* 2013;13:734–43.
- 69 Macedo LG, Hum A, Kuleba L, *et al.* Physical therapy interventions for degenerative lumbar spinal stenosis: a systematic review. *Phys Ther* 2013;93:1646–60.
- 70 Ammendolia C, Schneider M, Williams K, *et al.* The physical and psychological impact of neurogenic claudication: the patients' perspectives. *J Can Chiropr Assoc* 2017;61:18–31.
- 71 Friedly J, Chan L, Deyo R. Increases in lumbosacral injections in the Medicare population: 1994 to 2001. *Spine* 2007;32:1754–60.
- 72 Friedly J, Nishio I, Bishop MJ, *et al.* The relationship between repeated epidural steroid injections and subsequent opioid use and lumbar surgery. *Arch Phys Med Rehabil* 2008;89:1011–5.
- 73 Bresnahan BW, Rundell SD, Dagadakis MC, *et al.* A systematic review to assess comparative effectiveness studies in epidural steroid injections for lumbar spinal stenosis and to estimate reimbursement amounts. *Pm R* 2013;5:705–14.
- 74 Chou R, Hashimoto R, Friedly J, *et al.* Epidural corticosteroid injections for radiculopathy and spinal stenosis: a systematic review and meta-analysis. *Ann Intern Med* 2015;163:373–81.
- 75 Liu K, Liu P, Liu R, *et al.* Steroid for epidural injection in spinal stenosis: a systematic review and meta-analysis. *Drug Des Devel Ther* 2015;9:707–16.
- 76 Deyo RA, Mirza SK, Martin BI, *et al.* Trends, major medical complications, and charges associated with surgery for lumbar spinal stenosis in older adults. *JAMA* 2010;303:1259–65.
- 77 Machado GC, Maher CG, Ferreira PH, *et al.* Trends, complications, and costs for hospital admission and surgery for lumbar spinal stenosis. *Spine* 2017;42:1737–43.
- 78 Machado GC, Ferreira PH, Yoo RI, *et al.* Surgical options for lumbar spinal stenosis. *Cochrane Database Syst Rev* 2016;11:CD012421.
- 79 Zaina F, Tomkins-Lane C, Carragee E, *et al.* Surgical versus nonsurgical treatment for lumbar spinal stenosis. *Spine* 2016;41:E857–68.
- 80 Anderson DB, Mobbs RJ, Eyles J, *et al.* Barriers to participation in a placebo-surgical trial for lumbar spinal stenosis. *Heliyon* 2019;5:e01683.
- 81 Anderson DB, Ferreira ML, Harris IA, *et al.* Success, surgery for spinal stenosis: protocol of a randomised, placebo-controlled trial. *BMJ Open* 2019;9:e024944.
- 82 Enke O, New HA, New CH, *et al.* Anticonvulsants in the treatment of low back pain and lumbar radicular pain: a systematic review and meta-analysis. *CMAJ* 2018;190:E786–93.
- 83 Qaseem A, Wilt TJ, McLean RM, *et al.* Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2017;166:514–30.
- 84 Shanthanna H, Gilron I, Rajarathinam M, *et al.* Benefits and safety of gabapentinoids in chronic low back pain: a systematic review and meta-analysis of randomized controlled trials. *PLoS Med* 2017;14:e1002369.
- 85 Kim KH, Kim T-H, Lee BR, *et al.* Acupuncture for lumbar spinal stenosis: a systematic review and meta-analysis. *Complement Ther Med* 2013;21:535–56.
- 86 Bussi res AE, Stewart G, Al-Zoubi F, *et al.* Spinal manipulative therapy and other conservative treatments for low back pain: a guideline from the Canadian chiropractic guideline initiative. *J Manipulative Physiol Ther* 2018;41:265–93.
- 87 Schulz KF, Altman DG, Moher D, *et al.* Consort 2010 statement: updated guidelines for reporting parallel group randomised trials. *J Clin Epidemiol* 2010;63:834–40.
- 88 Chou R, Baisden J, Carragee EJ, *et al.* Surgery for low back pain: a review of the evidence for an American pain Society clinical practice guideline. *Spine* 2009;34:1094–109.
- 89 Kovacs FM, Urr tia G, Alarc n JD. Surgery versus conservative treatment for symptomatic lumbar spinal stenosis: a systematic review of randomized controlled trials. *Spine* 2011;36:E1335–51.
- 90 Suri P, Rainville J, Kalichman L, *et al.* Does this older adult with lower extremity pain have the clinical syndrome of lumbar spinal stenosis? *JAMA* 2010;304:2628–36.
- 91 Anderson SF, Kelley K, Maxwell SE. Sample-size planning for more accurate statistical power: a method adjusting sample effect sizes for publication bias and uncertainty. *Psychol Sci* 2017;28:1547–62.
- 92 Carlesso CPS, Smith C, Ammendolia C. Responsiveness of outcome measures in non-surgical patients with lumbar spinal stenosis a secondary analysis from a randomized controlled trial. *Spine* 2020.
- 93 Wertli MM, Buletti FC, Held U, *et al.* A comparison between different outcome measures based on "meaningful important differences" in patients with lumbar spinal stenosis. *Eur Spine J* 2017;26:450–61.
- 94 Ostelo RWJG, de Vet HCW. Clinically important outcomes in low back pain. *Best Pract Res Clin Rheumatol* 2005;19:593–607.
- 95 Boden SD, Davis DO, Dina TS, *et al.* Abnormal magnetic-resonance scans of the lumbar spine in asymptomatic subjects. A prospective investigation. *J Bone Joint Surg Am* 1990;72:403–8.
- 96 Haig AJ, Tong HC, Yamakawa KS, *et al.* Spinal stenosis, back pain, or no symptoms at all? A masked study comparing radiologic and electrodiagnostic diagnoses to the clinical impression. *Arch Phys Med Rehabil* 2006;87:897–903.
- 97 Genevay S, Courvoisier DS, Konstantinou K, *et al.* Clinical classification criteria for neurogenic claudication caused by lumbar spinal stenosis. The N-CLASS criteria. *Spine J* 2018;18:941–7.
- 98 Weinstein JN, Lurie JD, Tosteson TD, *et al.* Surgical compared with nonoperative treatment for lumbar degenerative spondylolisthesis. four-year results in the spine patient outcomes research trial (sport) randomized and observational cohorts. *J Bone Joint Surg Am* 2009;91:1295–304.
- 99 Weinstein JN, Tosteson TD, Lurie JD, *et al.* Surgical versus nonoperative treatment for lumbar spinal stenosis four-year results of the spine patient outcomes research trial. *Spine* 2010;35:1329–38.



## Nonoperative treatment for lumbar spinal stenosis – 22 July 2020 update

Database: Ovid MEDLINE: Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Ovid MEDLINE® Daily and Ovid MEDLINE® <1946-Present>

Search Strategy:

- 
- 1 randomized controlled trial.pt. (509927)
  - 2 controlled clinical trial.pt. (93770)
  - 3 Pragmatic clinical trial.pt. (1444)
  - 4 random\*.ti,ab. (1145458)
  - 5 placebo.ab,ti. (215288)
  - 6 drug therapy.fs. (2221199)
  - 7 trial.ab,ti. (599425)
  - 8 groups.ab,ti. (2097678)
  - 9 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 (5031369)
  - 10 (animals not (humans and animals)).sh. (4686362)
  - 11 9 not 10 (4375594)
  - 12 exp Constriction, Pathologic/ (30449)
  - 13 limit 12 to yr="1976 - 1982" (1906)
  - 14 exp Lumbar Vertebrae/ (52505)
  - 15 limit 14 to yr="1966 - 1982" (4472)
  - 16 exp Spinal Canal/ (7519)
  - 17 limit 16 to yr="1966 - 1982" (1172)
  - 18 exp Spinal Diseases/ (123399)
  - 19 limit 18 to yr="1966 - 1982" (18365)
  - 20 exp Spinal Stenosis/ (6116)
  - 21 spinal stenosis.ti,ab. (5088)
  - 22 (lumbar adj5 stenosis).ti,ab. (4268)
  - 23 (spin\* adj5 stenosis).ti,ab. (6620)
  - 24 neurogenic claudication.ti,ab. (633)
  - 25 exp Spinal Osteophytosis/ (4018)
  - 26 exp Spondylosis/ (7484)
  - 27 (lumb\* adj5 spondyl\*).ti,ab. (2886)
  - 28 exp Cauda Equina/ (3250)
  - 29 lumbar radicular pain.ti,ab. (218)
  - 30 13 or 15 or 17 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 (44520)
  - 31 9 and 30 (6508)
  - 32 limit 31 to ed=20190920-20200731 (275)
  - 33 limit 31 to yr=2019-2020 (545)
  - 34 32 or 33 (583)

Database: Embase Classic+Embase <1947 to 2020 July 21>

Search Strategy:

- 
- 1 Randomized Controlled Trial/ (613507)
  - 2 exp Controlled clinical trial/ (800817)
  - 3 Controlled Study/ (7533843)
  - 4 Double Blind Procedure/ (176652)

5 Single Blind Procedure/ (39549)  
 6 crossover procedure/ (64054)  
 7 placebo/ (362923)  
 8 Randomization/ (87513)  
 9 random\*.ti,ab. (1563918)  
 10 placebo?.ti,ab. (314621)  
 11 allocat\*.ti,ab. (155448)  
 12 assign\*.ti,ab. (400691)  
 13 blind\*.ti,ab. (436413)  
 14 (cross-over or crossover).ti,ab. (107060)  
 15 (compare or compared or comparing or comparison or comparative).ti,ab. (6802913)  
 16 (controlled adj7 (study or design or trial)).ti,ab. (355549)  
 17 ((singl\* or doubl\* or trebl\* or tripl\*) adj7 (blind\* or mask\*)).ti,ab. (250201)  
 18 trial.ti,ab. (878032)  
 19 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18  
 (12682849)  
 20 exp animals/ or exp invertebrate/ or animal experiment/ or animal model/ or animal tissue/ or  
 animal cell/ or nonhuman/ (29761121)  
 21 human/ or normal human/ or human cell/ (22533987)  
 22 20 and 21 (22470134)  
 23 20 not 22 (7290987)  
 24 19 not 23 (9386132)  
 25 exp vertebral canal stenosis/ (12543)  
 26 (spin\* adj5 stenosis).ti,ab. (9011)  
 27 (lumbar adj5 stenosis).ti,ab. (5728)  
 28 (neurogenic adj2 claudication).ti,ab. (1047)  
 29 (Spin\* adj2 Osteophytosis).ti,ab. (26)  
 30 exp cauda equina/ (4498)  
 31 lumbar radicular pain.ti,ab. (316)  
 32 (lumb\* adj5 spondyl\*).ti,ab. (4037)  
 33 exp spondylosis/ (9560)  
 34 spondylolisthesis/ (9419)  
 35 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 (36443)  
 36 24 and 35 (11296)  
 37 limit 36 to yr=2019-2020 (1405)  
 38 limit 36 to dd=20190920-20200731 (282)  
 39 37 or 38 (1426)

## CENTRAL via CRS Web

1 MESH DESCRIPTOR Spinal Stenosis EXPLODE ALL AND CENTRAL:TARGET 423  
 2 (spin\* NEAR5 stenosis) AND CENTRAL:TARGET 1189  
 3 lumb\* NEAR5 stenosis AND CENTRAL:TARGET 871  
 4 neurogenic claudication AND CENTRAL:TARGET 168  
 5 MESH DESCRIPTOR Spinal Osteophytosis EXPLODE ALL AND CENTRAL:TARGET 86  
 6 MESH DESCRIPTOR Spondylosis EXPLODE ALL AND CENTRAL:TARGET 374  
 7 lumb\* NEAR5 spondyl\* AND CENTRAL:TARGET 400  
 8 MESH DESCRIPTOR Cauda Equina EXPLODE ALL AND CENTRAL:TARGET 15  
 9 lumbar radicular pain AND CENTRAL:TARGET 93

10 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 AND CENTRAL:TARGET 1932  
 11 2019:YR AND CENTRAL:TARGET 105034  
 12 2020:YR AND CENTRAL:TARGET 30634  
 13 #11 OR #12 135668  
 14 #13 AND #10 209

## CINAHL

S43 S41 OR S42 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 242  
 S42 S40 AND EM 20190919-20200731 Search modes - Boolean/Phrase Interface - EBSCOhost  
 Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 192  
 S41 S40 Limiters - Published Date: 20190901-20200731  
 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 161  
 S40 S28 AND S39 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 3,036  
 S39 S29 OR S30 OR S31 OR S32 OR S33 OR S34 OR S35 OR S36 OR S37 OR S38 Search modes -  
 Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 6,262  
 S38 lumb\* W5 spondyl\* Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 796  
 S37 MH "Spondylolysis" Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 486  
 S36 MH "Spondylolisthesis" Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 1,438  
 S35 "lumbar radicular pain" Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 125  
 S34 MH "Cauda Equina" Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 368  
 S33 MH "Spinal Osteophytosis" Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search

Database - CINAHL Plus with Full Text 310  
 S32 "neurogenic claudication" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 243  
 S31 lumb\* W5 stenosis Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 1,768  
 S30 spin\* W5 stenosis Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 3,656  
 S29 MH "Spinal Stenosis" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 2,741  
 S28 S26 NOT S27 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 2,433,818  
 S27 MH "Animals" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 87,894  
 S26 S7 OR S12 OR S19 OR S25 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 2,461,016  
 S25 S20 OR S21 OR S22 OR S23 OR S24 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 1,686,740  
 S24 volunteer\* Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 52,797  
 S23 prospectiv\* Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 525,699  
 S22 control\* Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 1,275,002  
 S21 followup stud\* Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 203  
 S20 follow-up stud\* Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 12,011  
 S19 S13 OR S14 OR S15 OR S16 OR S17 OR S18 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases



Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 1,539,358  
S18 MH "Prospective Studies+" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 444,171  
S17 MH "Evaluation Research+" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 248,871  
S16 MH "Comparative Studies" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 331,705  
S15 latin square Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 248  
S14 MH "Study Design+" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 1,351,924  
S13 MH "Random Sample" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 34,389  
S12 S8 OR S9 OR S10 OR S11 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 431,064  
S11 random\* Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 414,911  
S10 placebo\* Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 66,332  
S9 MH "Placebos" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 12,827  
S8 MH "Placebo Effect" Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 2,282  
S7 S1 OR S2 OR S3 OR S4 OR S5 OR S6 Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search  
Database - CINAHL Plus with Full Text 404,557  
S6 triple-blind Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases

Search Screen - Advanced Search

Database - CINAHL Plus with Full Text 379  
 S5 single blind Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 15,679  
 S4 double blind Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 58,644  
 S3 clinical W3 trial Search modes - Boolean/Phrase Interface - EBSCOhost Research Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 250,481  
 S2 "randomi?ed controlled trial\*" Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 169,418  
 S1 MH "Clinical Trials+" Search modes - Boolean/Phrase Interface - EBSCOhost Research  
 Databases  
 Search Screen - Advanced Search  
 Database - CINAHL Plus with Full Text 303,246

PEDro

Yield: 12

Abstract and title: stenosis

AND

Body part: lumbar spine, sacroiliac joint or pelvis

AND

Method: clinical trial

Year: Since 2012

ICL

S1	, Peer Review only, Publication Type:Clinical Trial	85	2020-07-22 10:21:40
S2	, Peer Review only, Publication Type:Controlled Clinical Trial	9	2020-07-22
10:21:48			
S3	, Peer Review only, Publication Type:Randomized Controlled Trial	288	2020-
07-22 10:22:09			
S4	All Fields:random* OR All Fields:placebo* OR All Fields:sham, Peer Review only	1029	
2020-07-22 10:28:19			
S5	All Fields:"clinical trial" OR All Fields:"controlled trial", Peer Review only	481	
2020-07-22 10:28:38			
S6	All Fields:versus OR All Fields:vs., Peer Review only	196	2020-07-22 10:28:57
S7	All Fields:double-blind OR All Fields:"double-blind", Peer Review only	107	2020-
07-22 10:29:22			
S8	All Fields:single-blind OR All Fields:"single-blind", Peer Review only	456	2020-
07-22 10:29:33			
S9	, Peer Review only, Publication Type:Clinical Trial OR , Peer Review only, Publication		
Type:Controlled Clinical Trial OR , Peer Review only, Publication Type:Randomized Controlled Trial OR All			
Fields:random* OR All Fields:placebo* OR All Fields:sham, Peer Review only OR All Fields:"clinical trial"			
OR All Fields:"controlled trial", Peer Review only OR All Fields:versus OR All Fields:vs., Peer Review only			

OR All Fields:double-blind OR All Fields:"double-blind", Peer Review only OR All Fields:single-blind OR All Fields:"single-blind", Peer Review only 1584 2020-07-22 10:29:49

S10 Subject:"spinal stenosis" OR All Fields:"spinal stenosis", Peer Review only 83 2020-07-22 10:30:56

S11 Subject:"Spinal Osteophytosis" OR Subject:"Spondylosis" OR Subject:"Spondylolisthesis", Peer Review only 83 2020-07-22 10:31:27

S12 Subject:"Cauda equina" OR All Fields:"lumbar radicular pain", Peer Review only 14 2020-07-22 10:32:34

S13 Subject:"spinal stenosis" OR All Fields:"spinal stenosis", Peer Review only OR Subject:"Spinal Osteophytosis" OR Subject:"Spondylosis" OR Subject:"Spondylolisthesis", Peer Review only OR Subject:"Cauda equina" OR All Fields:"lumbar radicular pain", Peer Review only 172 2020-07-22 10:32:47

S14 , Peer Review only, Publication Type:Clinical Trial OR , Peer Review only, Publication Type:Controlled Clinical Trial OR , Peer Review only, Publication Type:Randomized Controlled Trial OR All Fields:random\* OR All Fields:placebo\* OR All Fields:sham, Peer Review only OR All Fields:"clinical trial" OR All Fields:"controlled trial", Peer Review only OR All Fields:versus OR All Fields:vs., Peer Review only OR All Fields:double-blind OR All Fields:"double-blind", Peer Review only OR All Fields:single-blind OR All Fields:"single-blind", Peer Review only AND Subject:"spinal stenosis" OR All Fields:"spinal stenosis", Peer Review only OR Subject:"Spinal Osteophytosis" OR Subject:"Spondylosis" OR Subject:"Spondylolisthesis", Peer Review only OR Subject:"Cauda equina" OR All Fields:"lumbar radicular pain", Peer Review only 26 2020-07-22 10:32:57

S15 , Year: from 2019 to 2020, Peer Review only 325 2020-07-22 10:33:21

S16 , Peer Review only, Publication Type:Clinical Trial OR , Peer Review only, Publication Type:Controlled Clinical Trial OR , Peer Review only, Publication Type:Randomized Controlled Trial OR All Fields:random\* OR All Fields:placebo\* OR All Fields:sham, Peer Review only OR All Fields:"clinical trial" OR All Fields:"controlled trial", Peer Review only OR All Fields:versus OR All Fields:vs., Peer Review only OR All Fields:double-blind OR All Fields:"double-blind", Peer Review only OR All Fields:single-blind OR All Fields:"single-blind", Peer Review only AND Subject:"spinal stenosis" OR All Fields:"spinal stenosis", Peer Review only OR Subject:"Spinal Osteophytosis" OR Subject:"Spondylosis" OR Subject:"Spondylolisthesis", Peer Review only OR Subject:"Cauda equina" OR All Fields:"lumbar radicular pain", Peer Review only AND , Year: from 2019 to 2020, Peer Review only

**Supplemental Table 2. Non operative interventions for neurogenic claudication due to lumbar spinal stenosis: A summary of GRADE assessment and outcomes (60 comparisons)**

						Walking ability/pain/function/quality of life measures				GRADE
Studies	Risk of Bias	Consistency	Directness	Precision	Selective Reporting	Immediate up to 1w	Short-term >1w - 3m	Intermediate 3m – 1yr	Long term ≥1yr	
<b>Calcitonin</b>										
<b>Calcitonin injection vs. placebo injection</b>										
Eskola 1992	High	No No	Yes Yes	No No	Yes		= TWT = VAS	= TWT = VAS	= TWT = VAS	+000 +000
Porter 1983	High	No	Yes	No	Yes		? Distance walked	? Distance walked		+000
Porter 1988	High	No No	Yes Yes	No No	Yes		= Distance walked = VAS			+000 +000
<b>Calcitonin nasal spray vs. placebo injection</b>										
Podichetty 2004	High	No No No No	Yes Yes Yes Yes	No No No No	Yes		= Distance walked = Time walked = SF-36 = VAS			+000 +000 +000 +000
Tafazal 2007	High	No No No No No	Yes Yes Yes Yes Yes	No No No No No	No		= Shuttle walk = VAS leg = VAS back = ODI = Global			+000 +000 +000 +000 +000
<b>Calcitonin nasal spray plus physical therapy vs. paracetamol plus physical therapy</b>										
Sahin 2009	High	No No No	Yes Yes Yes	No No No	No		= Distance walked = VAS = RMDI			+000 +000 +000
<b>Oral Medication</b>										
<b>Oral prostaglandin vs. Etodlac (NSAID)</b>										
Matsudaira 2009	Low	No No No No No	Yes Yes Yes Yes Yes	No No No No No	Yes		> Distance walked # ? SF-36 = LBP > Leg pain > Global #			++00 +000 ++00 ++00 ++00
<b>Methylcobalamin (vit B12) plus conservative care vs. conservative care</b>										
Waikukul 2000	High	No	Yes	No	No			> Distance walked #	> Distance walked #	+000



Gabapentin plus physical therapy, corset & NSAIDS vs. placebo plus physical therapy, corset & NSAIDS										
Yaksi 2007	High	No No No	Yes Yes Yes	No No No	No		= VAS	> Distance walked > VAS	> Distance walked # > VAS #	+000 +000
Pregabalin vs. active placebo										
Markman 2015	High	No No No No	Yes Yes Yes Yes	No No No No	No	= NPS rest/final = Distance walked = Recovery time = Global < RMDQ				+000 +000 +000 +000 +000
Gabapentin plus conservative vs. conservative plus botulinum										
Park 2017	High	No No No	Yes Yes Yes	No No No	No		= NPS (Back/leg) = ODI = Global			0000 0000 0000
Oxymorphone hydrochloride vs. placebo										
Markman 2015 - 2	High	No No No No No No	Yes Yes Yes Yes Yes Yes	No No No No No No	No	= NPS rest/final = Distance walked = Recovery Time = ZCQ (s) = ZCQ (f) = Global				0000 0000 0000 0000 0000 0000
Propoxyphene/acetaminophen vs. placebo										
Markham 2015 – 2	High	No No No No No No	Yes Yes Yes Yes Yes Yes	No No No No No No	No	= NPS rest/final = Distance walked = Recovery Time = ZCQ (s) < ZCQ (f) # = Global				0000 0000 0000 0000 0000 0000
Oxymorphone hydrochloride vs. propoxyphene/acetaminophen										
Markham 2015 - 2	High	No No No No No No	Yes Yes Yes Yes Yes Yes	No No No No No No	No	= NPS rest/final = Distance walked = Recovery Time = ZCQ (s) > ZCQ (f) # = Global				0000 0000 0000 0000 0000 0000
Oral corticoid vs. placebo										

Rodrigues 2014	High	No No No No	Yes Yes Yes Yes	No No No No	No		= SF-36 = RMDQ = 6 min walk < VAS #			0000 0000 0000 0000
<b>Rehabilitation Therapy and Multimodal Care</b>										
Exercise plus ultrasound vs. exercise plus sham ultrasound										
Goren 2010	low	No No No No	Yes Yes Yes Yes	No No No No	No		= TWT = VAS back = VAS leg # = ODI			++00 ++00 ++00 ++00
Exercise plus ultrasound vs. no treatment										
Goren 2010	Low	No No No No	Yes Yes Yes Yes	No No No No	No		= TWT = VAS back > VAS leg # > ODI			++00 ++00 ++00 ++00
Exercise plus sham ultrasound vs. no treatment										
Goren 2010	Low	No No No No	Yes Yes Yes Yes	No No No No	No		= TWT = VAS back > VAS leg # > ODI #			++00 ++00 ++00 ++00
In-patient physical therapy vs. home exercise program plus oral diclofenac										
Koc 2009	High	No No No No	Yes Yes Yes Yes	No No No No	Yes		= TWT = VAS = RMDI = NHP	= TWT = VAS = RMDI = HNP		+000 +000 +000 +000
Unweighted treadmill walking plus exercise vs. cycling plus exercise										
Pua 2007	Low	No No No No No	Yes Yes Yes Yes Yes	No No No No No	No		= Distance walked = ODI = RMDI = VAS = Global			++00 ++00 ++00 ++00 ++00
Manual therapy, exercise and unweighted treadmill vs. flexion exercise, walking and sham ultrasound										
Whitman 2006	High	No No No No	Yes Yes Yes Yes	No No No No	No		= TWT > Global # = ODI = NPRS			+000 +000 +000 +000
Supervised physical therapy vs home exercises										

Minetama 2019	High	No	Yes	No	No		> ZCQ (F) #			+000
		No	Yes	No			>ZCQ (S) #			+000
		No	Yes	No			> Distance walked #			+000
		No	Yes	No			> NPS (leg)			
		No	Yes	No			> SF-36 PF			+000
		No	Yes	No			> SF-36 BP			+000
		No	Yes	No			= Daily Steps			+000
Manual therapy & exercise vs medical care										
Schneider 2019	Low	No	Yes	Yes	No		> ZCQ #	= ZCQ		+++0
		No	Yes	Yes			= SPWT	= SPWT		+++0
		No	Yes	Yes			= PA	= PA		+++0
Manual therapy & exercise vs. community exercise										
Schneider 2019	Low	No	Yes	Yes	No		> ZCQ #	= ZCQ		+++0
		No	Yes	Yes			= SPWT	= SPWT		+++0
		No	Yes	Yes			= PA	= PA		+++0
Community exercise vs. medical care										
Schneider 2019	Low	No	Yes	Yes	No		= ZCQ	= ZCQ		+++0
		No	Yes	Yes			= SPWT	= SPWT		+++0
		No	Yes	Yes			= PA	= PA		+++0
Comprehensive therapy and exercise vs. self-directed exercise										
Ammendolia 2018	Low	No	Yes	Yes	No	> SPWT #	> SPWT #	> SPWT #	> SPWT #	+++0
		No	Yes	Yes		> 30% SPWT	> 30% SPWT	> 30% SPWT	>30% SPWT	+++0
		No	Yes	Yes		> 50% SPWT	= 50% SPWT	= 50% SPWT	> 50% SPWT	+++0
		No	Yes	Yes		> ZCQ (s)	= ZCQ (s)	= ZCQ (s)	> ZCQ (f) #	+++0
		No	Yes	Yes		= ZCQ (f)	= ZCQ (f)	= ZCQ (f)	> ZCQ (s) +	+++0
		No	Yes	Yes		= ODI	= ODI	> ODI (walk)	ZCQ (f)	+++0
		No	Yes	Yes		> NPS (back) #	= NPS (back)	= NPS (back)	= ODI	+++0
		No		Yes		= NPS (leg)	= NPS (leg)	= NPS (leg)	= NPS (back)	+++0
		No				= SF-36 BP	= SF-36 BP	= SF-36 BP	> SF-36 BP #	+++0
		No				= SF-36 PF	> SF-36 PF #	= SF-36 PF	>SF-36 PF #	+++0
Standard exercise vs. isokinetic exercises										
Oğuz 2013	High	No	Yes	No	Yes	= VAS	= VAS	= VAS		0000
		No	Yes	No		= ODI	= ODI	= ODI		0000
		No	Yes	No		= TWT	= TWT	= TWT		0000
Standard exercise vs. unloaded exercise										
Oğuz 2013	High	No	Yes	No	Yes	< VAS	< VAS	= VAS		0000
		No	Yes	No		< ODI	= ODI	= ODI		0000
		No	Yes	No		= TWT	= TWT	= TWT		0000

Isokinetic exercises vs. unloaded exercises										
Oğuz 2013	High	No No No	Yes Yes Yes	No No No	Yes	< VAS < ODI < TWT #	= VAS < ODI < TWT	= VAS = ODI = TWT		0000 0000 0000
Aquatic physical therapy exercise vs. physical therapy										
Homayouni 2015	High	No No	Yes Yes	No No	Yes	> VAS # > Distance walked	= VAS = Distance walked			0000 0000
Pre-surgical exercise program vs. routine preoperative hospital management										
Marchand 2019	High	No No	Yes Yes	No No	Yes	> NPS (leg) # > Duration walked #	= NPS (leg) = Duration walked	= NPS (leg) = Duration walked		0000 0000
Gang-Chuk Tang (herbal concoction), daily Mokuri Chuna therapy, daily acupuncture, physician consultation vs. oral aceclofenac, epidural steroid injection, physical therapy										
Kim 2019	Low	No No No No	Yes Yes Yes Yes	No No No No	Yes		= VAS (leg) = VAS (back) > OCS > Distance walked	= VAS (leg) > VAS (back) # = OCS > Distance walked		+000 +000 +000 +000
Mokhuri Chuna, acupuncture, and physician consultation vs. oral aceclofenac, epidural steroid injection, physical therapy										
Kim 2019	Low	No No No No	Yes Yes Yes Yes	No No No No	Yes	>VAS (low back)#	= VAS (leg) = VAS (back) = OCS = Distance walked	> VAS (leg) # > VAS (back) # = OCS = Distance walked		+000 +000 +000 +000
Spinal Manipulation										
Lumbar spinal manipulation vs. waiting										
Passmore 2017	High	No No	Yes Yes	No No	No	= NPS (Back) = NPS (Leg)				0000 0000
Acupuncture										
Acupuncture with usual care vs. usual care										



Kim 2016	High	No	Yes	No	No		6 weeks: = ODI = SF-36 BP = SF-36 PF = LBP = Leg pain = Distance walked 3 months: = ODI = SF-36 BP = SF-36 PF = LBP = Leg pain = Distance walked			0000 0000 0000 0000 0000 0000 0000 0000 0000 0000
Acupuncture vs. sham acupuncture										
Qin 2020	Low	No	Yes	No	No	> RMDQ > NRS (back) # > NRS (leg) # > SSS-S # > SSS-F # = SPWT	> RMDQ > NRS (back) # > NRS (leg) # > SSS-S # > SSS-F # = SPWT	> RMDQ > NRS (back) > NRS (leg) # > SSS-S # > SSS-F # = SPWT		++00 ++00 ++00 ++00 ++00 ++00
Epidural Injection										
Translaminar epidural steroid injections vs. placebo injections										
Cuckler 1985	High	No	Yes	No	No	= Global			=global	+000
Translaminar epidural steroids plus epidural block vs. placebo injections										
Fukusaki 1988	High	No	Yes	No	No	> Distance walked #	= Distance walked			+000
Translaminar epidural steroids plus epidural block vs. epidural block injections										
Fukusaki 1988	High	No	Yes	No	No	= Distance walked	= Distance walked			+000
Translaminar epidural block vs. placebo										
Fukusaki 1988	High	No	Yes	No	No	> Distance walked #	= Distance walked			+000
Intralaminar epidural steroid plus epidural block vs. home exercise program plus oral diclofenac										
Koc 2009	High	No	Yes	No	Yes		= TWT > VAS # > RMDI #	= TWT = VAS = RMDI		+000 +000 +000

		No	Yes	No	Yes		> NHP	= HNP		+000
Intralaminar epidural steroid plus epidural block vs. in-patient physical therapy										
Koc 2009	High	No	Yes	No	Yes		= TWT	= TWT		+000
		No	Yes	No	Yes		= VAS	= VAS		+000
		No	Yes	No	Yes		= RMDI	= RMDI		+000
		No	Yes	No	Yes		= NHP	= HNP		+000
Caudal epidural steroids vs. placebo injections										
Zahaar 1991	High	No	Yes	No	No	= Global			= Global	+000
Mild lumbar decompression vs. epidural steroid injection										
Brown 2012	High	No	Yes	No	No		= VAS			0000
		No	Yes	No			= ODI			0000
		No	Yes	No			= ZCQ			0000
		No	Yes	No			12 weeks:			
							= VAS			0000
		No	Yes	No			= ODI			0000
		No	Yes	No			= ZCQ			0000
Lidocaine vs. glucocorticoid–lidocaine										
Friedly 2014, 2017	Low	No	Yes	Yes	No		3 weeks:	12 weeks:	12 months:	
		No	Yes	Yes			< RMDQ	= RMDQ	= RMDQ	+++0
							< NPS (leg)	= NPS (leg)	= NPS (leg)	+++0
		No	Yes	Yes			6 weeks:	6 months:		
		No	Yes	Yes			= RMDQ	= RMDQ		+++0
							= NPS (leg)	= NPS (leg)		+++0
Makris 2016	Low	No	Yes	No	Yes		Makris 2016			
							3 weeks:			
		No	Yes	No	Yes		< RMDQ using SIP			0000
		No	Yes	No	Yes		Weights			
							< RMDQ Patient-			0000
							Prioritized			
		No	Yes	No	Yes		(LESSER)			
		No	Yes	No	Yes		6 weeks:			
							< RMDQ using SIP			
							Weights			0000
		No	Yes	No	Yes		= RMDQ Patient-			
							Prioritized			0000
							(LESSER)			

Lidocaine spinal injection vs. saline spinal injection										
Song 2016	High	No No	Yes Yes	No No	No		1 month: = VAS = FRI 3 months: = VAS = FRI			0000 0000 0000 0000
Fluoroscopically guided lumbar ILESIS at the level of maximal stenosis vs. two intervertebral levels cephalad										
Milburn 2014	High	No No No	Yes Yes Yes	No No No	No	1 week: > NPS (walking) # > RMDQ #	4 weeks: > NPS (walking) # > RMDQ 12 weeks: = NPS (walking) > RMDQ			0000 0000 0000 0000
Epidural steroid injection (ESI) Vs. ESI & physiotherapy										
Hammerich 2019	High	No No No No No	Yes Yes Yes Yes Yes	No No No No No	No		= ODI = NPS > SF-36 ER # > SF-36 EWB > SF-36 GH	= ODI = NPS # = SF-36 ER = SF-36 EWB = SF-36 GH	= ODI > NPS # = SF-36 ER = SF-36 EWB = SF-36 GH	0000 0000 0000 0000 0000
Interlaminar vs. transforaminal epidural steroid injection										
Sencan 2020	High	No No No No	Yes Yes Yes Yes	No No No No	Yes	= NPS	3 weeks: = NPS = ODI > BDS = Distance walked	3 months: > NPS = ODI > BDS > Distance walked #		0000 0000 0000 0000 0000 0000 0000 0000
TNF alpha inhibitor (Etanercept) vs. steroid injection										
Wei 2020	Low	No No No	Yes Yes Yes	No No No		> VAS #	1, 3 months: > VAS # > ODI #	6 months: > VAS # > ODI #		++00 ++00 ++00
TNF alpha inhibitor (Etanercept) vs. lidocaine										

Wei 2020	Low	No No No	Yes Yes Yes	No No No		> VAS #	1, 3 months: > VAS # > ODI #	6 months: > VAS # > ODI #		++00 ++00 ++00
Steroid vs. lidocaine injection										
Wei 2020	Low	No No No	Yes Yes Yes	No No No		= VAS	1, 3 months: = VAS = ODI	6 months: = VAS = ODI		++00 ++00 ++00
Percutaneous Epidural Adhesiolysis										
Balloon-less catheter (Racz) vs. inflatable balloon catheter (ZiNeu)										
Karm 2018	High	No No No  No No No	Yes Yes Yes  Yes Yes Yes	No No No  No No No	No		1 month: = NPS (back) = NPS (leg) = ODI 3 months: = NPS (back) = NPS (leg) = ODI	6 months: < NPS (back) # < NPS (leg) # < ODI		0000 0000 0000  0000 0000 0000
Surgery vs. Physical Therapy										
Interspinous spacer (X Stop) vs. non operative care										
Zucherman 2004, 2005, Hsu 2006	High	No No	Yes Yes	No No	No		> ZCQ(S)# > ZCQ(F)# > SF-36 PF > SF-36 BP > SF-36 GH > SF-36 ER	> ZCQ(S)# > ZCQ(F)# > SF-36 PF > SF-36 BP > SF-36 GH > SF-36 ER	> ZCQ(S)# > ZCQ(F)# > SF-36 PF# > SF-36 BP# > SF-36 GH > SF-36 ER#	+000 +000 +000 +000 +000 +000
Laminectomy +/- fusion vs. non operative care for degenerative spondylolisthesis										

Weinstein 2007, 2009 Abdu 2018	High	No No No No No	Yes Yes Yes Yes	No No No No	No		= SF-36 BP, PF = ODI = LBPBS = LPBI = SBS	= SF-36 BP, PF = ODI = LBPBS = LPBI = SBS	2 years: = SF-36 BP, PF = ODI = LBPBS = LPBI = SBS 4 years: = SF-36 BP, PF = ODI = LBPBS = LPBI = SBS 8 years: = SF-36 BP, PF = ODI = LBPBS = LPBI = SBS	+000 +000 +000 +000 +000 +000 +000 +000 +000 +000 +000 +000 +000 +000
Laminectomy +/- fusion vs. non operative care										
Amundsen 2000	High	No No	Yes Yes	No No	No		?* Pain severity	?* Global	?* Pain severity ? Global	+000 +000
Malmivaara 2007 N= 94	Low	No No No No No	Yes Yes Yes Yes	No No No No	No			= TWT = SW > VAS leg walk # > VAS LB walk # > ODI	= TWT = SW > VAS leg walk # > VAS LB walk # > ODI	++00 ++00 ++00 ++00 ++00
Weinstein 2008, 2010, Lurie 2015	High	No No No No No	Yes Yes Yes Yes Yes	No No No No	No		= SF-36 BP = SF-36 PF = LBPBS = LPBI = SBS = ODI	= SF-36 BP = SF-36 PF = LBPBS = LPBI = SBS = ODI	2 years: > SF-36 BP ** # = SF-36 PF = LBPBS = LPBI = SBS = ODI 4 years: =SF-36 BP ** = SF-36 PF = LBPBS = LPBI	+000 +000 +000 +000 +000 +000 +000 +000 +000 +000 +000 +000



									= SBS 8 years: = SF-36 BP = SF-36 PF = ODI = Stenosis Index	+000 +000 +000 +000
Laminectomy, facet resection, neuroforaminotomy vs. physical therapy										
Delitto 2015	High	No No	Yes Yes	No No	No				2 years: = SF-36 = ODI	+000 +000

> favours intervention (first comparison), < favours control (second comparison), = no difference between intervention and control groups, TWT= Treadmill Walking Test, VAS= Visual Analog Scale for Pain Intensity, RMDI= Roland-Morris Back Disability Index, NHP= Nottingham Health Profile, Global= Patient Perceived Improvement, SR= Selective Reporting, ODI= Oswestry Back Disability Index, ?= insufficient data, LBP= Low back Pain Severity Scale, Leg pain= Leg Pain Severity Scale, ? SF-36=No data on overall score, improvement in some subscales, NPRS= Numeric Pain Rating Scale, SF-36 BP= SF-36 Bodily Pain Subscale, SF-36- PF= SF-36 Physical Function Subscale, SF-36 ER= SF-36 emotional role subscale, SF-36 EWB= SF-36 emotional well-being subscale, SF-36 GH= SF-36 General health subscale, LBPBS= Low Back Pain Bothersome Scale, LPBI= Leg Pain Bothersome Index, SBS= Stenosis Bothersome Scale, SW= Subjective Walking, VAS leg= Visual Analog Scale for Leg Pain, VAS LB= Visual Analog Scale for Low Back Pain, VAS leg walking= Visual Analog Scale for Leg pain while walking, SIP= sickness index profile, BDS= Beck Depression Score, LESSER= Lumbar Epidural Steroid Injection for Spinal Stenosis Extended Research, PA= Physical Activity, FRI= Functional Rating Index, TWT= Total Walking Time, SSS= Spinal Stenosis Questionnaire, ?\*= no between group statistical comparisons, \*\*= SF-36 BP significantly better at 2 years but not 4 years.

GRADE evidence; +000= Very low GRADE evidence, ++00= Low GRADE, +++0= Moderate GRADE evidence, ++++= High GRADE evidence

# between group difference meeting the MCID. The MCID used were:  $\geq 1.25$  points for back pain and  $\geq 1.5$  points for leg pain on 0 to 100-point Visual Analogue Scale (VAS) and 0 to 10-point Numerical Rating Scale (NRS) for back pain (58),  $\geq 5$  points on 0- to 24-point Roland-Morris Disability Questionnaire (RMDQ) (59),  $\geq 8$  points for conservative treatment and  $\geq 12$  points for surgery on 0- to 100-points for Oswestry Disability Index (ODI) (60),  $\geq 0.1$  points for the functional component and 0.36 points for symptom component of the Zurich Claudication Questionnaire (ZCQ) (58),  $\geq 0.38$  points for combined symptoms and functional scores of the ZCQ (92),  $\geq 30\%$  between-group difference for walking distance, global improvement and SF36 subscales (61).

Supplemental Table 1. Characteristics of included studies

Study	Participants and Settings	Interventions	Outcomes/Follow-up	Results (Group 1 is reference group)
<b>Calcitonin</b>				
Eskola 1992	39 subjects with an average of 6 years of pain, average age of 56.6 years of age, 20 males and 19 females.  <b>Setting:</b> Orthopaedic hospital in Finland.	1) 100IU Calcitonin injection every other day for 4 weeks (n=20)  2) Placebo treatment (Miacalcic Sandoz 100IU) every other day for 4 weeks (n=19)	1) VAS 2) Treadmill test 3) Coping with ADLs 4) Digitest 5) Ergojump 6) Blood tests  <b>Follow-up:</b> 1, 3, 4, 6 and 12 months	<b>Between group WMD and 95% CI</b> <b>Pain (VAS) (mm):</b> -0.050 (-0.053 to -0.047) <b>Walking distance (meters):</b> -18.5 (-240.37 to 203.37)  <b>Adverse events:</b> The calcitonin injection group reported minor nausea and rash in 89% of the subjects.
Podichetty 2004	55 subjects with an average age of 68.5 years and an average of 36.2 weeks of the condition in the intervention group and 29.8 weeks in the placebo group, 33 males and 22 females.  <b>Setting:</b> Spinal center in the United States	1) 400 IU intranasal calcitonin daily for 6 weeks followed by open label 6-week extension (n=36)  2) Placebo nasal spray daily for 6 weeks, followed by open label 6-week extension, during which all patients received 400IU calcitonin (n=19)	1) VAS 2) Walking capacity 3) ODI 4) Stenosis specific questionnaire 5) Satisfaction with pain levels, functional status, and treatment received 6) SF-36 7) Symptom diary  <b>Follow-up:</b> 12 weeks	<b>Between group MD, 95% CI, p values</b> <b>12 weeks:</b> <b>Pain VAS (mm):</b> 0.5 (-0.85 to 1.93); p=0.44, <b>Walking time (seconds):</b> 42.2 (-86.9 to 170.4); p=0.51 <b>Walking distance (feet):</b> 163.3 (-311.16 to 637.84); p=0.049 <b>SF-36 MCS:</b> -4.22 (-10.41 to 1.97); p=0.18 <b>SF-36 PCS:</b> 0.43 (-3.73 to 4.59); p=0.84
Porter	41 subjects with	1) 100 IU salmon calcitonin injection	1) Walking chart	Insufficient data provided to calculate mean difference in

1983	10 in a double blind RCT crossover, 37 males and 4 females with mean age of 55.4 years.  <b>Setting:</b> Infirmary in England	four times per week, sometimes with Maxalon for nausea (n=5)  2) Matching placebo (n=5)  Only responders randomized	and ability to walk more than 1 mile  2) ODI  <b>Follow-up:</b> 10 weeks	walking distance or ODI among the 10 patients enrolled in RCT.  <b>Adverse events:</b> The calcitonin injection group reported minor nausea and rash in 40% of the subjects.
Porter 1988	42 subjects, 35 male, 7 female, average age of 53.6 years in 20 subjects and 56.7 years in 22 subjects, median duration of back pain reported was 11 years for 19 subjects, and 14 years for 22 subjects. Median duration of claudication was 1.25 years for 20 subjects and 4.5 years for 22 subjects.  <b>Setting:</b> Infirmary in England	1) 100 IU of salmon calcitonin injected subcutaneously 4 times per week for 8 weeks (n=20)  2) 1 ml of saline injected 4 times per week for 8 weeks (n=22)	1) VAS 2) Claudication threshold 3) 3 level mobility assessment 4) Analgesic requirements 5) 3 level sleep disturbance 6) Treatment success defined as 100% improvement in walking distance and able to walk 800 m.  <b>Follow-up:</b> 4 and 8 weeks	<b>Difference in median score from baseline between groups</b> <b>Pain score (VAS) (mm):</b> 4 weeks: -9 8 weeks: -5.5 <b>Walking distance until symptoms onset (meters):</b> 4 weeks: -14 8 weeks: 42 <b>Walking distance until pain prevents walking (meters):</b> 4 weeks: -41 8 weeks: -99  No significant between group differences. No p values or 95% CI provided.
Sahin 2009	45 subjects 31 males and 14 females, average	1) 200 IU intranasal calcitonin daily for 8 weeks (n=23)	1) VAS 2) Walking capacity	<b>Percent change between groups:</b> <b>8 weeks:</b> <b>VAS at rest:</b> 4.7%, p>0.05

	<p>ages of 57.65 years in calcitonin group and 54.45 years in paracetamol group.</p> <p><b>Setting:</b> Physical and Rehabilitation Medicine Department in Turkey</p>	<p>2) Up to 1500mg of paracetamol daily for 8 weeks (n=22)</p> <p>Both groups took part in a physical therapy and exercise program 5 times per week for 15 sessions.</p>	<p>3) RMDI</p> <p>4) Ranges of motion</p> <p><b>Follow-up:</b> 8 weeks</p>	<p><b>VAS with motion:</b> -7.9%, P&gt;0.05</p> <p><b>Roland Morris:</b> 8.2%, p&gt;0.05</p> <p><b>Walking distance:</b> -15.4%, p&gt;0.05</p>
Tafazal 2007	<p>40 subjects, 30 males, 10 females, average of 67 years in the intervention group and 70.2 years in the placebo group, average of 38.7 months with symptoms in the calcitonin group and 30.9 months in the placebo group.</p> <p><b>Setting:</b> University hospital in England</p>	<p>1) Placebo nasal spray NaCl for 4 weeks (n=20)</p> <p>2) 200 IU nasal salmon calcitonin for 4 weeks (n=20)</p>	<p>1) VAS</p> <p>2) Shuttle walking test</p> <p>3) 4-point subjective outcome of overall assessment (excellent, good, fair, poor)</p> <p>4) ODI</p> <p>5) Modified Somatic Perception Questionnaire</p> <p>6) Modified Zung Depression Score</p> <p><b>Follow-up:</b> Baseline, 4, 10, 16 weeks</p>	<p><b>4 weeks: Between group MD 95% CI</b></p> <p><b>ODI:</b> -0.7 (1.7 to -3.5)</p> <p><b>LBOS:</b> -3.0 (-0.6 to -4.7)</p> <p><b>VAS leg (mm):</b> -10 (-4.0 to -13)</p> <p><b>VAS back (mm):</b> -6.0 (-6 to -12)</p> <p><b>Shuttle walk distance (m):</b> -13 (-7 to -35)</p> <p><b>16 weeks: between group MD, p values</b></p> <p><b>ODI:</b> 0.1, p=0.44;</p> <p><b>LBOS:</b> 0.7, p=0.93;</p> <p><b>VAS leg (mm):</b> -4, p=0.66;</p> <p><b>VAS back (mm):</b> 16, p=0.03;</p> <p><b>Shuttle walking distance (m):</b> -11, p=0.39</p>

Oral Medication				
Matsudaira 2009	79 subjects, 24 males and 24 females, with an average age of 69.6 years in the Limaprost group and 72.2 in the Etodolac group.  <b>Setting:</b> Orthopaedic surgery in a medical faculty in Japan	1) Oral prostaglandin E1 derivative (15 g Limaprost) 3 times daily for 8 weeks (n=39)  2) 400 mg of etodolac (NSAID) twice daily for 8 weeks (n=40)	1) SF-36 2) Verbal pain rating scales 3) Walking distance 4) LBP severity 5) Leg pain severity 6) Leg numbness severity 7) Treatment satisfaction  <b>Follow-up:</b> 8 weeks	<b>SF-36 subscales MD, p values</b> 8 weeks: <b>physical function:</b> 9.4, p=0.01, <b>role physical:</b> 13.7, p=0.03, <b>bodily pain:</b> 15.5, p<0.01; <b>General health:</b> 6.6, p=0.08; <b>vitality:</b> 11.3, p=0.02; <b>social functioning:</b> 8.0, p=0.17; <b>role emotional:</b> 10.2, p=0.07; <b>mental health:</b> 12.2, p<0.01.  <b>Secondary outcomes</b> not provided in a way that MD can be extracted: 8 weeks: <b>low back pain:</b> p=0.77; leg pain p=0.08; <b>Leg numbness:</b> p<0.01; <b>walking distance</b> p<0.01; <b>patient subjective improvement</b> p<0.01; <b>patient satisfaction</b> p<0.01 all in favor of limaprost  <b>Adverse events:</b> 5% of subjects in both groups reported gastrointestinal upset.
Waikakul 2000	152 subjects, 68 males and 84 females with an average age of 66.8 years. 44 of the subjects had symptoms for less than one month, 98 had symptoms for more than one month.  <b>Setting:</b> Hospital in Thailand	1) Conservative treatment consisting of education, activity modification, exercise and physical therapy. NSAIDs, muscle relaxants, and analgesics as necessary. Vitamin B1, B6, and B12 3 times per day (n=82)  2) Conservative treatment plus Methlcobalin ESAI, 1.5mg per day in 3 divided doses after meals for 6 months (n=70)	1) Presence of pain on spinal motion 2) Claudication distance 3) Medication intake (NSAIDs, muscle relaxants, and steroids)  <b>Follow-up:</b> every month for two years	<b>Walking distance</b> <b>Percent able to walk &gt; 1000 meters</b> 6 mo: 71.3% vs. 88.6%, p< 0.05 12 mo: 81.3% vs. 97.1%, p < 0.05 18mo: 83.8% vs. 97.1% p < 0.05  <b>Adverse events:</b> There were no reported adverse effects in subjects in methylocabalin group
Yaksi 2007	55 subjects, 22 males, 33 females, average age of 50.8 years.  <b>Setting:</b> Hospital	1) 900 mg of gabapentin per day increased weekly by 300 mg to a maximum of 2400 mg (n=28)  2) Placebo (n=27)	1) VAS – low back and leg pain during movement 2) Walking distance	<b>Between group difference, p values</b> <b>Pain (VAS) (mm) no raw data</b> 3 <sup>rd</sup> mo 3.4 vs. 1.9, p =0.039 4 <sup>th</sup> mo 4.1 vs.2.0, p =0.006  <b>Walking Ability, no raw data</b>



	department of physical medicine and rehabilitation in Turkey	Both groups received physical therapy exercises, a lumbosacral corset with steel bracing and NSAID treatments	3) Presence or absence of motor and/or sensory deficits  <b>Follow-up:</b> 15 days, 1, 2, 3, 4 months	Grp 1: longer walking distance at end of 2 <sup>nd</sup> mo (p < 0.05), 3 <sup>rd</sup> mo (p < 0.05) and 4 <sup>th</sup> mo (p < 0.005)  <b>Adverse events:</b> some subjects randomized to the gabapentin group (no data specified) experienced mild to moderate drowsiness and/or dizziness.
Markman 2015	29 participants, 20 males, 9 females, Eligible subjects were older than 50 years (mean 70.1 years) with at least one level of radiographically confirmed lumbar spinal stenosis and symptoms of neurogenic claudication for at least 3 months.  <b>Setting:</b> Hospital in Rochester, New York	1) Pregabalin group (n=14)  2) Active placebo (Diphenhydramine) (n=15)  Cross over study after 7 day wash out period. Pregabalin was started at 75 mg PO twice daily or diphenhydramine, 6.25 mg) and increased on day 4 to 150 mg PO twice daily (12.5 mg diphenhydramine) for 7 days. Pregabalin was decreased to 75 mg PO twice daily (6.25 mg diphenhydramine) on day 11 for 3 days of tapering.	1) NRS - time to first moderate pain symptom during a 15-minute treadmill test (Tfirst) (NRS - greater than 4)  <b>Follow-up:</b> day 10 of intervention period	<b>Between group MD, 95% CI, p values</b> <b>Treadmill testing pain at rest (NRS)</b> 0.29 (0.41 to 0.98): p=0.40 <b>Treadmill testing final pain (NRS)</b> 0.25 (-0.44 to 0.94): p=0.46 <b>Treadmill testing distance walked (m)</b> -24.06 (-75.63 to 27.52): p=0.35 <b>Treadmill testing recovery time (min)</b> -0.79 (-1.86 to 0.28): p=0.14 <b>Treadmill testing patient global assessment of pain</b> -0.08 (-0.45 to 0.29): p=0.67 <b>Treadmill testing RMDQ</b> 1.50 (0.38 to 2.62): p=0.01  <b>Adverse events:</b> Complications were reported in 64% of subjects in group 1, the most common being dizziness, compared to 35% in group 2.
Park 2017	45 subjects, 21 in GPN Group (17 female, 4 males, mean age 66.1±10.5), and 24 in BTX group (15 female and 9 males, mean age	1) Conservative treatments plus gabapentin (group GPN): Gabapentin 300 to 1200mg/d - titrated to patient characteristics, comorbidities, and reported side effects (n=21)  2) Conservative treatments plus BTX	3) NRS - back/leg pain intensity  4) Cramp frequency (no./wk)  5) Cramp severity (0-4)	No statistically significant difference between groups and lack of reporting of quantitative data  <b>Adverse events:</b> Five patients (20.8%) in group 2 reported mild to moderate pain at injection sites for a few days.

	66.2±8.2)  <b>Setting:</b> Outpatient department for interventional pain management in Korea	injection (group BTX): The BTX (botulinum toxin type A [Nabota]) dose was 100U in 5mL of 0.9% saline injected into the gastrocnemius medialis and lateralis. (n=24)  Conservative treatments: education, exercise, analgesic medication, injection therapy including epidural injections, and physical therapy	criteria) 6) Insomnia severity – (ISI 0-28) 7) ODI 8) Patient global impression of change  <b>Follow-up:</b> 2 weeks, 1 and 3 months.	
Markman 2015 - 2	24 participants, 12 males and 12 females, (mean age 72 years) LSS by imaging with symptoms of neurogenic claudication  <b>Setting:</b> Translational Pain Research Center at a University in Rochester, New York	1) Oxymorphone hydrochloride (Opana IR, 5 mg) (n=8) 2) Propoxyphene/acetaminophen (Darvocet, 100 mg/650 mg) (n=8) 3) Placebo: 3 separate visits (random order with at least 3 day washout periods) (n=8)	1) NRS (at rest) 2) NRS (final pain rating) 3) AUC 4) Distance walked (m) 5) Recovery time (min) 6) ZCQ 7) Patient global assessment of pain 8) RMDQ 9) ODI  <b>Follow-up:</b> Study was prematurely terminated	<b>Between group MD, 95% CI, p values</b> <b>Treadmill testing pain at rest (NRS)</b> Grp 1 vs Grp 3: -0.04 (-0.72 to 0.65): p=0.89 Grp 2 vs Grp 3: -0.27 (-0.95 to 0.41): p=0.32 Grp 1 vs Grp 2: 0.23 (-0.45 to 0.92): p=0.40 <b>Treadmill testing final pain (NRS)</b> Grp 1 vs Grp 3: 0.2 (-0.74 to 1.14): p=0.60 Grp 2 vs Grp 3: 0.53 (-0.40 to 1.46): p=0.16 Grp 1 vs Grp 2: -0.33 (-1.26 to 0.61): p=0.39 <b>Treadmill testing distance walked (m)</b> Grp 1 vs Grp 3: -12.41 (-63.01 to 38.20): p=0.54 Grp 2 vs Grp 3: -23.41 (-73.60 to 26.79): p=0.25 Grp 1 vs Grp 2: 11 (-39.53 to 61.54): p=0.59 <b>SSSQ symptom severity score</b> Grp 1 vs Grp 3: -0.03 (-0.19 to 0.13): p=0.61 Grp 2 vs Grp 3: 0.01 (-0.15 to 0.17): p=0.85 Grp 1 vs Grp 2: -0.04 (-0.20 to 0.11): p=0.49 <b>SSSQ physical function score</b> Grp 1 vs Grp 3: 0.04 (-0.16 to 0.09): p=0.47 Grp 2 vs Grp 3: 0.11 (-0.01 to 0.23): p=0.03 Grp 1 vs Grp 2: -0.15 (-0.27 to -0.02): p=0.01 <b>Patient global assessment of pain</b> Grp 1 vs Grp 3: -0.03 (-0.52 to 0.47): p=0.90 Grp 2 vs Grp 3: 0.13 (-0.36 to 0.61): p=0.52

				Grp 1 vs Grp 2: -0.15 (-0.64 to 0.34): p=0.44  <b>The study was prematurely terminated because of the removal of propoxyphene/acetaminophen from the US market.</b>
Rodrigues 2014	61 patients with lumbar canal stenosis (50–75 years; canal area < 100 mm <sup>2</sup> at L3/L4, L4/L5, and/or L5/S1 on MRI; and claudication within 100 m). 31 in the corticoid group (mean age 58.23 (6.38), and 30 in the placebo group (mean age 58.33 (6.19))  <b>Setting:</b> Hospital in São Paulo, Brazil	1) Oral corticoid group received 1 mg/kg of oral corticoids daily, with a dose reduction of one-third per week for 3 weeks (n=31)  2) Control group was administered placebo for the same period (n=30)	1) SF-36 2) RMDQ 3) 6-min walk test 4) VAS 5) Likert scale  <b>Follow-up:</b> 3, 6 and 12 weeks	<b>Between group comparison VAS (6 weeks)</b> Corticoid vs Placebo: 1.53 p=0.02 (in favour of placebo)
<b>Rehabilitation Therapy and Multimodal Care</b>				
Goren 2010	45 subjects, 13 males, 32 females, average ages in groups of 57.4, 49.13, and 53.06. 7 subjects with pain duration of 3-6 months, 7 with pain duration of 6-12 months, and	1) Stretching and strengthening exercises for lumbar, abdominal, leg muscles as well as low intensity cycling exercises were given as therapeutic exercises. Ultrasound was applied with 1mHz, 1.5W/cm <sup>2</sup> intensity, in continuous mode on the back muscle for 10 minutes (n=17)  2) Same as group 1 with Ultrasound on off- mode (n=17)	1) VAS (out of 10) 2) Treadmill test at 3 km/h for maximum of 15 minutes or 750m. 3) ODI 4) Analgesic consumption 5) Physiatrist	<b>Pain (VAS) (mm) within group MD</b> 3 weeks: Grp 1: -2.2 for <b>back pain</b> ; -1.47 for <b>leg pain</b> Grp 2: -1.94 for <b>back pain</b> ; -2.47 for <b>leg pain</b> Grp 3: 0.40 for <b>back pain</b> ; 0.54 for <b>leg pain</b>  Between groups differences Leg pain: Grp 1 > Grp 3 (p<0.01), Grp 2 > Grp 3 (p<0.01)  <b>Walking Ability (within group MD)</b> 3 weeks: Grp 1: 94.30 seconds

	31 with pain duration of greater than 12 months.  <b>Setting:</b> Rehabilitation center in Turkey	3) No exercise-no treatment (n=16)	assessment  <b>Follow-up:</b> End of 3-week treatment period only	Grp 2: 114.94 seconds Grp 3: -66.10 seconds No significant change between groups  <b>Disability (ODI) (within group MD)</b> 3 weeks: Grp 1: -3.94 Grp 2: -7.8 Grp 3: -3.6  ODI between groups differences Grp 1> Grp 3 (p<0.05), Grp 2> Grp 3 (p<0.05)
Koc 2009	29 subjects, 21 male, 8 female, average ages of 62.6, 61.1, and 53.1 years in the three groups, average pain duration of 5.7 years, 5.0 years, and 5.7 years in the three groups.  <b>Setting:</b> Medical school department of physical medicine and rehabilitation in Turkey	1) Conservative inpatient physical therapy program 5 days a week for 2 weeks. PT included applications of ultrasound 1.5 W/cm <sup>2</sup> for 10min, hot pack for 20min, and TENS for 20min to the lumbar region (n=13)  2) Lumbar epidural steroid injections, 10 ml of solution containing 60mg of triamcinolon acetone (1.5 mL), 15 mg of 0.5% bupivacain hydrochloride (3 mL), and 5.5 mL of physiologic saline (0.9%NaCl) was injected in 3.5minutes. (n=10)  3) Control group (n=10)  All patients included were trained to pursue a home-based therapeutic exercise program performed twice daily for a period of 6 months, and oral diclofenac sodium 75mg was administered to all patients twice daily for 2 weeks	1) VAS 2) Treadmill walk test 3) Nottingham Health Profile 4) RMDI 5) Functional testing including finger to floor distance, sit-to-stand, and a weight carrying test  <b>Follow-up:</b> 2 weeks, 1, 3 and 6 months	No raw data provided. No significant between group differences for all outcomes and follow-ups except:  <b>Pain (VAS)</b> 2 weeks: Grp 2 less pain than Grp 3 p= 0.008  <b>Disability (RMDI)</b> 2 weeks: Grp 2 less disability than Grp 3 p= 0.007  <b>Quality of Life (Nottingham Health Profile) (no data provided)</b> Grp 2 had significantly higher improvement than Grp 3 at 2 weeks in mobility subgroup scores.  <b>Adverse events:</b> 1 subject reported angina pectoralis and 1 reported gastric complaints (group not specified).
Pua 2007	68 subjects, 35 males, 33	1) Unweighted treadmill training: Weeks 1 and 2, participants walked	1) VAS for pain over past	<b>Pain (VAS) (mm) MD and 95% CI</b> 6 weeks: 2 ( -5 to 10)

	<p>females, average age of 58 years, 12 week median pain duration</p> <p><b>Setting:</b> Hospital in Singapore</p>	<p>with a relatively pain-free gait which translated to 30–40% of body weight. In weeks 3 to 6, participants were encouraged to walk at a moderate intensity. The duration of each treadmill session was limited by participant tolerance or to a maximum of 30 minutes. 2x per week for 6 weeks = 12 sessions (n=33)</p> <p>2) Cycling on upright bicycle: During weeks 1 and 2, participants cycled at their comfortable pace at 50 to 60 rpm. Participants were instructed to assume a flexed posture. In weeks 3 to 6, participants were encouraged to exercise at a moderate intensity and the duration of each cycling session was limited by participant tolerance or to a maximum of 30 minutes. 2x per week for 6 weeks for 12 sessions (n=35)</p>	<p>week</p> <p>2) Patient perceived benefit on a 6-point scale</p> <p>3) ODI</p> <p>4) RMDI</p> <p>5) Walking ability</p> <p><b>Follow-up:</b> 3 and 6 weeks</p>	<p><b>Disability (ODI), OR, 95% CI</b> 6 weeks: OR 1.10 (0.41 to 2.98)</p> <p><b>Patient perceived benefit, OR, 95% CI</b> 6 weeks: OR 0.50 (0.17 to 1.48)</p> <p><b>Walking ability (≥800 m), OR, 95% CI</b> 6 weeks: OR 1.14 (0.44 to 2.94)</p> <p><b>Adverse events:</b> 1 subject in treadmill group reported increase in pain.</p>
Whitman 2006	<p>58 subjects, 31 males, 27 female, 29 (group 1) with an average age of 70 years, 29 (group 2) with an average age of 68.9, median low back pain duration of 108 months in Group 1's 29 subjects and 60 months in Group 2's 29</p>	<p>1) Flexion Exercise and Walking Group: 45-60 minutes twice per week for 6 weeks. Lumbar flexion exercises along with self-pace treadmill walking program, and sub-therapeutic ultrasound. The duration of each treadmill session was based on that patient's tolerance on that specific day and could extend up to 45 minutes. (n=29)</p> <p>2) Manual Therapy, Exercise and Walking Group: 45-60 minutes twice per week for 6 weeks - Manual</p>	<p>1) Global Rating of Change (15-point scale)</p> <p>2) NPRS for lower limb</p> <p>3) Walking Tolerance test</p> <p>4) ODI</p> <p>5) Medication consumption</p> <p>6) Satisfaction subscale of the Spinal</p>	<p><b>Patient Global Assessment (somewhat better or greater)</b> 6 weeks: 41% vs. 79% p&lt;0.01 1 year: 21% vs. 38% p&gt;0.05</p> <p><b>Number needed to treat for benefit for perceived recovery and 95% CI</b> 6 weeks: 2.6 (1.8 to 7.8) 1 year: 4.8 (-2.3 to 21.3) long term: 4.4 (- 2.1 to 22.7)</p> <p><b>Pain (NPRS lower extremity) Within group MD, 95% CI</b> 6 weeks: 1.1 (0.2 to 2.0) vs. 1.5 (0.5 to 2.5) 1 year: 1.2 (0.4 to 1.9 vs. 1.0 (-0.2 to 2.2);</p>



	<p>subjects, lower extremity median pain duration of 48 months in Group 1's 29 subjects and 24 months in Group 2's 29 subjects.</p> <p><b>Setting:</b> University in the United States</p>	<p>physical therapy (thrust and non thrust) to the thoracic and lumbar spine, pelvis, and lower extremities and specific exercises at discretion based on the underlying impairments. Patients received specific exercises to address impairments in mobility, strength, and/or coordination. Exercises were performed in the clinic and as part of a home exercise program. Patients also underwent a bodyweight supported treadmill ambulation program using a cable and trunk harness system to unload a specific amount of weight from the patient while the patient walks as comfortably as possible on a treadmill (n=29).</p>	<p>Stenosis Scale</p> <p>7) Additional use of health care resources</p> <p><b>Follow-up:</b> 6 weeks, 1 year, long term mail survey (averaging 29 months)</p>	<p>Long term: 1.8 (0.6 to 3.0) vs. 2.0 (0.7 to 3.4) Between group MD not statistically significant at any follow-up period</p> <p><b>Walking Ability (improvement in meters) within group MD, 95% CI</b> 6 weeks: 176.5 (-9.5 to 362.4) vs. 339.7 (218.4 to 461) 1 year: 130.4 (-55.3 to 316.2) vs. 209.8 (67.5 to 352.1) Between group improvement not statistically significant at any follow-up</p> <p><b>Disability (ODI) within group MD</b> 6 weeks: 6.55 (1.87 to 11.23) vs. 10.48 (6.5 to 14.4) 1 year: 5.03 (1.71 to 8.35) vs. 7.14 (1.5 to 12.8) Between group differences not statistically significant at any follow-up</p>
Minetama 2019	<p>86 patients, 39 men and 47 women, average age 72.7 years 43 patients (20 men and 23 women, average age 72.3 years to the PT group 43 patients (19 men and 24 women, average age 73.2 years) to the HE group. Duration symptoms 20 months</p>	<p>1) Physical therapy + home exercise program (n=43) 2) Home exercise (HE) program alone (n=43)</p> <p>Supervised physical therapy twice a week for 6 weeks, including manual therapy, individually tailored stretching and strengthening exercises, cycling, and body weight-supported treadmill walking. The manual therapy included manipulation, stretching, and massaging of the thoracic and lumbar spine, pelvis, and lower extremities. The individually tailored muscle exercises included those for the trunk (eg, abdominal planks, side bridge, and/or back extension) and lower</p>	<p>1) ZCQ 2) Satisfaction 3) SPWT (m) 4) NRS 5) JOABPEQ-acquired points 6) SF-36 7) HADS 8) PCS 9) PASS-20 10) TSK-11 11) Daily steps</p> <p><b>Follow-up:</b> 6 weeks</p>	<p><b>Between group MD, 95% CI</b> <b>ZCQ - Symptom severity</b> -0.4 (-0.6 to -0.2): statistically significant <b>ZCQ - Physical function</b> -0.4 (-0.6 to -0.2): statistically significant <b>SPWT (m)</b> 455.9 (308.5 to 603.2): statistically significant <b>NRS - Leg pain</b> -1.4 (-2.5 to -0.3): statistically significant <b>SF-36 - Physical functioning</b> 9.2 (2.1 to 16.3): statistically significant <b>SF-36 - Bodily pain</b> 10.4 (3.3 to 17.5): statistically significant <b>Daily steps</b> 723.4 (199.1 to 1,283.5): statistically significant</p>

	<p><b>Setting:</b> Spine care center at a university hospital in Japan</p>	<p>extremities (eg, unloading hip and/or knee exercise with ankle weight and/or standing squats). The typical dosage for strengthening exercises was a total of 2 to 3 sets with 10 repetitions, each of 6-second contraction. The typical duration of stretching was three repetitions of 30 seconds.</p> <p>All patients in both groups were asked to take a daily walk that did not exacerbate their lower extremity symptoms using a pedometer and walking diary and to perform a HE program consisting of lumbar flexion exercises including three 30-second bouts of both single and double knee-to-chest exercises, ten 6-second bouts of trunk raises and bridging in the supine position, and a 4-point kneeling exercise at least twice daily.</p>		
Schneider 2019	<p>259 subjects, 122 males and 137 women with an average age of 72.4, 68 patients had symptoms for less than 6 months, 191 had symptoms for greater than 6 months</p> <p><b>Setting:</b> Outpatient research clinic in Pittsburgh</p>	<p>1) Medical care (MC) (n=88)</p> <p>2) Group exercise (GE) (n=84)</p> <p>3) Manual therapy + exercise (MTE) (n=87)</p> <p><b>Medical Care:</b> 3 visits to a physical medicine physician over 6 weeks. Primarily prescription of oral medications in any combination of nonnarcotic analgesics, anticonvulsants, antidepressants. Optional referral for epidural steroid injections if inadequate pain relief by oral medication, severe neurogenic claudication, and/or patient preference.</p>	<p>1) SSS</p> <p>2) SPWT</p> <p>3) Physical Activity</p> <p><b>Follow-up:</b> 2 and 6 months</p>	<p><b>Between group MD, 95% CI</b></p> <p><b>SSS (2 months)</b></p> <p>GE vs MC: 0.4 (-1.3 to 2.1)</p> <p>MTE vs MC: -2.0 (-3.6 to -0.4)</p> <p>MTE vs GE: -2.4 (-4.1 to -0.8)</p> <p><b>SPWT (2 months)</b></p> <p>GE vs MC: 79.9 (-74.5 to 234.5)</p> <p>MTE vs MC: 122.9 (-25.7 to 271.6)</p> <p>MTE vs GE: 43.0 (-111.8 to 197.9)</p> <p><b>Physical activity (2 months)</b></p> <p>GE vs MC: 28.7 (2.7 to 54.7)</p> <p>MTE vs MC: 20.4 (-4.5 to 45.3)</p> <p>MTE vs GE: -8.3 (-34.5 to 17.6)</p> <p><b>SSS (6 months)</b></p> <p>GE vs MC: -0.5 (-2.3 to 1.3)</p> <p>MTE vs MC: -1.1 (-2.8 to 0.6)</p> <p>MTE vs GE: -0.6 (-2.4 to 1.2)</p>

		<p>Physician rendered general guide and on gentle stretching and advice to stay active.</p> <p><b>Group Exercise:</b> Supervised exercise classes at 2 local senior community centers. 2x 45-min classes/week, 6 weeks. Taught by senior fitness instructors. Participants self-select level of exercise based on fitness level (easy to medium)</p> <p><b>Manual Therapy + Exercise:</b> 2x 45minute sessions per week, 6 weeks by either 2 chiropractors or 2 physiotherapists. Sessions included 3 interventions:</p> <ol style="list-style-type: none"> <li>1. Warm-up procedure on stationary bicycle</li> <li>2. Manual therapy procedures (lumbar distraction, hip, lumbar/sacroiliac joint and neural mobilizations)</li> <li>3. Individualized instruction in spinal stabilization exercises and home stretching</li> </ol> <p>Practitioner determined what muscles required stretch/strengthening and appropriate exercises added to program.</p>		<p><b>SPWT (6 months)</b>  GE vs MC: 86.5 (-75.7 to 248.8)  MTE vs MC: 73.8 (-84.1 to 231.7)  MTE vs GE: -12.7 (-175.6 to 150.1)</p> <p><b>Physical activity (6 months)</b>  GE vs MC: 21.3 (-6.9 to 49.4)  MTE vs MC: -2.9 (-30.1 to 24.3)  MTE vs GE: -24.2 (-52.5 to 4.0)</p> <p><b>Adverse events:</b> There were no reported serious adverse events in any group. There was a significantly greater rate of transient joint soreness associated with group 3 (49%) compared with group 2 (31%) and group 1 (6%).</p>
Ammendolia 2018	104 patients, 45 males and 59 females, 48 in comprehensive group and 51 in self-directed group, with an average age of 69.4	<ol style="list-style-type: none"> <li>1) Comprehensive (n=48)</li> <li>2) Self-directed (n=51)</li> </ol> <p><b>Comprehensive:</b> Chiropractor providing 2x/week of 15-20-minute treatment sessions over a 6-week period followed by a single (booster) session, 4 weeks later.</p>	<ol style="list-style-type: none"> <li>1) SPWT Distance</li> <li>2) Clinical Significance - 30% improvement in SPWT no. (%)</li> <li>3) Clinical</li> </ol>	<p><b>Between group MD, 95% CI, p values</b></p> <p><b>SPWT</b>  8 wks: 345.4 (150.0 to 540.7): p=0.00  3 mo: 304.1 (77.9 to 530.3): p=0.01  6 mo: 421.0 (181.4 to 660.6): p=0.00  12 mo: 473.2 (203.9 to 742.4): p=0.00</p> <p><b>30% improvement in SPWT</b>  8 wks: 24 (6-40): p=0.01  3 mo: 21 (4-38): p=0.02</p>

	<p>(comprehensive) and 71.7 (self-directed) neurogenic claudication &gt;3 months, imaging-confirmed canal narrowing, walk &gt;20m and not surgical candidates in next 12 months</p> <p><b>Setting:</b> Academic hospital outpatient clinic in Toronto</p>	<p>Education: Self-management strategies via cognitive behavioral approach. Body repositioning (pelvic tilt) when standing and walking.</p> <p>Exercises: Standardized set of exercises demonstrated gradually over 6 weeks and was a part of structured home exercise program. Cycling, muscle stretching, strengthening, conditioning for back and lower extremity fitness and to facilitate lumbar flexion</p> <p>Manual therapy: Spinal manipulation; joint, soft tissue and neural mobilization; lumbar flexion-distraction; and manual muscle stretching applied each visit. Participants received an instructional video and workbook and pedometer.</p> <p><b>Self-directed:</b> Instructional Video, workbook, pedometer and a single 15-to 30-minute training session with an experienced independent licensed chiropractor, independent of the comprehensive program, Training session: Describe 6-week program, review workbook, explain pedometer use and recording of weekly walking steps. Video and workbook: Educational information and the same exercise instruction and self-management strategies received by the comprehensive group</p>	<p>Significance - 50% improvement in SPWT no. (%)</p> <p>4) ZCQ-S 5) ZCQ-F 6) ZCQ-S + ZCQ-F 7) ODI 8) ODI walk 9) NRS Back 10) NRS Leg</p> <p><b>Follow-up:</b> 8 weeks, 3, 6, and 12 months</p>	<p>6 mo: 19 (2-35): p=0.02 12 mo: 22 (4-39): p=0.02 <b>50% improvement in SPWT</b> 8 wks: 26 (8-42): p=0.01 3 mo: 19 (-1.0 to 36): p=0.06 6 mo: 17 (-2 to 35): p=0.09 12 mo: 24 (5-40): p=0.01 <b>ZCQS</b> 8 wks: -0.19 (-0.37 to -0.02): p=0.03 3 mo: -0.15 (-0.37 to 0.08): p=0.19 6 mo: -0.02 (-0.22 to 0.19): p=0.87 12 mo: -0.22 (-0.47 to 0.02): p=0.07 <b>ZCQF</b> 8 wks: -0.02 (-0.22 to 0.17): p=0.81 3 mo: -0.18 (-0.39 to 0.03): p=0.09 6 mo: -0.11 (-0.33 to 0.11): p=0.34 12 mo: -0.27 (-0.49 to 0.04): p=0.02 <b>ZCQS+ZCQF</b> 8 wks: -0.24 (-0.56 to 0.07): p=0.13 3 mo: -0.36 (-0.75 to 0.03): p=0.07 6 mo: -0.23 (-0.58 to 0.12): p=0.20 12 mo: -0.48 (-0.90 to -0.06): p=0.03 <b>ODI</b> 8 wks: -0.02 (-0.07 to 0.02): p=0.30 3 mo: -0.04 (-0.09 to 0.01): p=0.13 6 mo: -0.02 (-0.07 to 0.02): p=0.34 12 mo: -0.03 (-0.08 to 0.02): p=0.30 <b>ODI Walk</b> 8 wks: -0.2 (-0.6 to 0.1): p=0.14 3 mo: -0.4 (-0.9 to 0.03): p=0.07 6 mo: -0.9 (-1.3 to -0.4): p&lt;0.001 12 mo: -0.2 (-0.7 to 0.2): p=0.32 <b>NRS Back</b> 8 wks: -1.4 (-2.2 to -0.5): p=0.002 3 mo: -0.6 (-1.4 to 0.3): p=0.23 6 mo: -0.7 (-1.7 to 0.3): p=0.16 12 mo: -0.4 (-1.3 to 0.4): p=0.32</p>
--	---	--	--	--

				<p><b>NRS Leg</b>  8 wks: -0.7 (-1.5 to 0.1): p=0.09  3 mo: 0.05 (-0.85 to 0.96): p=0.91  6 mo: -0.9 (-1.9 to 0.003): p=0.58  12 mo: -0.5 (-1.6 to 0.6): p=0.37</p> <p><b>SF-36 Bodily Pain</b>  8 wks: 2.0 (-4.9 to 8.9): p=0.57  3 mo: -4.5 (-12.4 to 3.5): p=0.27  6 mo: -3.3 (-10.2 to 3.6): p=0.35  12 mo: 10 (2.1 to 17.9): p=0.013</p> <p><b>SF-36 Physical Function</b>  8 wks: 4.2 (-3.9 to 12.4): p=0.31  3 mo: 9.2 (1.1 to 17.3): p=0.027  6 mo: 5.8 (-2.1 to 13.6): p=0.15  12 mo: 8.2 (0.2 to 16.2): p=0.045</p> <p><b>Adverse events:</b> At 12 months, 0 participants out of 43 in group 1 and 2 out of 46 participants in group 2 experienced adverse events that were mostly attributed to a temporary increase in low back and/or leg pain.</p>
Oğuz 2013	<p>120 patients, 30 in group 1 with an average age of 57.1 years old, 30 in group 2 with an average age of 55.8 years old and group 3 with an average age of 57.4 years old, LSS symptoms, narrowing by MRI</p> <p><b>Setting:</b> University</p>	<p>1) Standard exercise group (n=30)  2) Isokinetic exercise program (n=30)  3) Unloading exercise group (n=60)</p> <p>All groups physician-guided (5x/week for 3 weeks) then at-home (3x/week)</p> <p><b>Standard Exercise:</b> 15 sessions of TENS, hot packs with home exercise instruction.</p> <p><b>Isokinetic exercise:</b> 20 minutes/day, 5 sessions/week for a total of 15 sessions with a physician. Isokinetic exercises:</p>	<p>1) VAS  2) ODI  3) Beck Depression Inventory</p> <p><b>Follow-up:</b> 4, 12 and 24 weeks</p>	<p><b>Between group MD, p value</b></p> <p><b>VAS</b>  After treatment:  Grp 1 vs Grp 2: 0.37, p&gt;0.05  Grp 1 vs Grp 3: 1.36, p&lt;0.05  Grp 2 vs Grp 3: 0.99, p&lt;0.05  4<sup>th</sup> week:  Grp 1 vs Grp 2: 1.43, p&gt;0.05  Grp 1 vs Grp 3: 1.17, p&lt;0.05  Grp 2 vs Grp 3: -0.26, p&gt;0.05  12<sup>th</sup> week:  Grp 1 vs Grp 2: 0.93, p&gt;0.05  Grp 1 vs Grp 3: 0.71, p&gt;0.05  Grp 2 vs Grp 3: -0.22, p&gt;0.05  24<sup>th</sup> week:  Grp 1 vs Grp 2: 1.08, p&gt;0.05</p>



	department of physical medicine and rehabilitation in Turkey	<p>rates of 60°/sec, 120°/sec, 180°/sec with 70° of body movement (50° flexion to 20° extension) Each session had 3 sets, each set had 5 repetitions at described velocity, with 20s rest between each set.</p> <p><b>Unloaded exercise:</b> 5 sessions of unloading exercise per week, for a total of 15 sessions with a physician. Walking with unloading exercise device: session 1-5 = 45% body weight, session 6-15 = 30% body weight. Treadmill walking at 1.2 km/hr for 20 minutes, or until pain due to neurogenic claudication was felt. Subjects advised to follow exercise program s at home at least 3x/week after discharge.</p>	<p>Grp 1 vs Grp 3: 0.46, p&gt;0.05 Grp 2 vs Grp 3: -0.62, p&gt;0.05 <b>ODI</b> After treatment: Grp 1 vs Grp 2: -0.8, p&gt;0.05 Grp 1 vs Grp 3: 1.8, p&lt;0.05 Grp 2 vs Grp 3: 2.6, p&lt;0.05 4<sup>th</sup> week: Grp 1 vs Grp 2: 1.5, p&gt;0.05 Grp 1 vs Grp 3: 2.6, p&gt;0.05 Grp 2 vs Grp 3: 1.1, p&lt;0.05 12<sup>th</sup> week: Grp 1 vs Grp 2: 1, p&gt;0.05 Grp 1 vs Grp 3: 1.3, p&gt;0.05 Grp 2 vs Grp 3: 0.3, p&gt;0.05 24<sup>th</sup> week: Grp 1 vs Grp 2: 0.4, p&gt;0.05 Grp 1 vs Grp 3: 0.5, p&gt;0.05 Grp 2 vs Grp 3: 0.1, p&gt;0.05 <b>Total Gait Duration</b> After treatment: Grp 1 vs Grp 2: 64.6, p&gt;0.05 Grp 1 vs Grp 3: -50.5, p&gt;0.05 Grp 2 vs Grp 3: -115.1, P&lt;0.05 4<sup>th</sup> week: Grp 1 vs Grp 2: 45.9, p&gt;0.05 Grp 1 vs Grp 3: -18.4, p&gt;0.05 Grp 2 vs Grp 3: -64.3, p&lt;0.05 12<sup>th</sup> week: Grp 1 vs Grp 2: 52.23 p&gt;0.05 Grp 1 vs Grp 3: -0.67 p&gt;0.05 Grp 2 vs Grp 3: -52.9 p&gt;0.05 24<sup>th</sup> week: Grp 1 vs Grp 2: 35.2, p&gt;0.05 Grp 1 vs Grp 3: 1.9, p&gt;0.05 Grp 2 vs Grp 3: -33.3, p&gt;0.05</p>
--	--	---	---

Homayouni 2015	<p>47 subjects, 23 male, 24 female, 24 in group one, mean age 55.56, 12 male, 12 female, 23 in group two, mean age 55.68, 11 male, 12 female</p> <p><b>Setting:</b> University-based pain clinics in Iran</p>	<p>1) Treatment in therapeutic pools with water temperature of 29–30 degrees Celsius. Every aquatic session started with warm up and ended with cool down, with duration of 10–15 min for each of them. Participants should have attended aquatic physical therapy sessions every other day for a total duration of 24 sessions. Each session included ambulation, side walking, chain walking, forward walking with kickboard, stretching of each muscle group including adductors, abductors, flexors and extensors of the hip, knee flexors and ankle plantar flexors and dorsiflexors. Other interventions were mini-squat, pelvic curl, pelvic tilt, and knee to chest, double knee lift, and deep-water exercise. (n=25)</p> <p>2) Passive modalities by physical therapists including continuous mode ultrasound (US) 1.5W/ cm<sup>2</sup> for 10 min and hot pack and trans-electrical nerve stimulation (TENS) for 20 min to the lumbar region. Also, the therapists instructed the patients in this group to perform trunk muscle endurance, William's and stretching exercises. The patients were treated using these passive modalities and were given exercises under supervision of physiotherapists for 10 sessions. They were instructed to perform the learned exercises 30 min</p>	<p>1) VAS 2) Walking ability</p> <p><b>Follow-up:</b> Immediately after therapy, 3 months</p>	<p><b>All between group comparisons</b> <b>Walking ability</b> Grp 1 &gt; Grp 2: p=0.02 <b>VAS</b> Grp 1 &gt; Grp 2 p=0.001</p>
----------------	---	--	---	---

		a day at home in the following weeks until the end of the eighth week. (n=25)		
Marchand 2019	40 participants, 17 females and 23 males, 20 in the intervention group with an average age of 66.7 years old and 20 in the control group with an average age of 71.5 years old, with history and diagnostic imaging of LSS  <b>Setting:</b> Regional hospital in Quebec	1) Exercise 3x week / 6 weeks prior to surgery (n=20)  2) Regular hospital preoperative management with back posture education (n=20)	1) NRS (Pain Intensity) 2) ROM (Active) 3) Muscle strength (N-m) 4) Walking capacity (seconds)  <b>Follow-up:</b> 3 and 6 months	<b>Between group MD</b> <b>NRS (leg)</b> Preoperative: -2.1, p<0.05 Postoperative: 1.1, p>0.05 3 months: 1.1, p>0.05 6 months: 0.3, p>0.05 <b>ROM (active)</b> Preoperative: 5, p<0.05 Postoperative: -6, p>0.05 <b>Muscle Strength</b> Preoperative: 45.7, p<0.001 Postoperative: 5.1, p>0.05 <b>Walking Duration</b> Preoperative: 90, p<0.05 Postoperative: -14.5, p>0.05
Kim 2019	34 subjects, mean age 64 (5.3), women 24 (66.7)  <b>Setting:</b> Hospital in Seoul, South Korea	1) MT1 group: 110 g of Gang-Chuk Tang was administered 3 times a day (Gang-Chuk Tang is an herbal concoction consisting of Eucommiae Cortex, Achyranthis Radix, Rhizoma Cibtii, Sorbus commixta, G. thunbergii, Saposhnikovia Radix, and Acanthopanacis Cortex in equal portions) Daily Mokhuri Chuna therapy (relaxation and mobilization of lumbar joint and back muscle) Daily acupuncture treatment on LI4, ST36, LV3, BL22, BL23, BL24, BL25, and Ashi points. Consultation on precautions related to daily	1) VAS for leg pain 2) VAS for low back pain 3) Oxford Claudication Scoring 4) Walking distance  <b>Follow-up:</b> 3 and 6 months	<b>All between group comparisons</b> <b>VAS leg pain (post treatment)</b> MT2 (28.82±27.46) vs CMT (51.82±25.34) groups: P=0.04 <b>VAS leg pain (6 months)</b> MT1 (48.91±23.08) vs CMT (72.27±16.72) groups: P=0.01 MT2 (42.36±21.29) vs CMT groups: P=0.003 <b>VAS low back pain (6 months):</b> MT2 (30.00±13.48) vs CMT (60.82±18.62) groups: P=0.001 <b>Oxford Claudication Scoring (3 months)</b> MT1 (18.75±6.52) vs CMT (25.82±6.24) groups: p=0.02 <b>Walking distance (3 months)</b> MT1 vs CMT: p=0.03 <b>Walking distance (6 months)</b> MT1 vs CMT: p=0.01

		<p>activity and stepwise walking training for the entire 4 weeks of therapy. (n=12)</p> <p>2) MT2 group: Mokhuri Chuna, acupuncture, and physician consultation were offered in the same manner and dosage as the MT1 group with the exception that all herbal medications were withheld. (n=11)</p> <p>3) CMT group: Oral analgesic therapy (aceclofenac 100 mg twice daily and eperisone hydrochloride 50 mg three times daily for 28 days) and three interlaminar epidural steroid injections (5 mg of dexamethasone per injection) at the level of the affected spinal region over a 4-week period were administered. Physiotherapy including heating pad, and transcutaneous electrical nerve simulator, and deep tissue heating therapy five times per week for 4 weeks. (n=11)</p>		The primary outcome of this pilot study was safety as measured by the type and incidence of adverse events (AEs).
<b>Spinal Manipulation</b>				
Passmore 2017	14 patients with degenerative LSS (n=14); Swiss Spinal Stenosis score of M=63.2, standard deviation [SD] = 15.9 (mean age 59.0 (10.6)), 7 in the SM group (4	<p>1) Spinal manipulation group: received bilateral high-velocity; low-amplitude spinal manipulation directed toward the lumbar region (by a licensed chiropractor with more than 10 years of clinical experience) (n=7)</p> <p>2) Non Intervention Group: Waited 5 minutes if they were assigned to the</p>	<p>1) Movement time</p> <p>2) NPS (Back)</p> <p>3) NPS (leg)</p> <p>4) ROM</p> <p><b>Follow-up:</b> Immediately after intervention</p>	<p>There was no significant difference between groups for all outcomes.</p> <p>1. Grp 1 vs. Grp 2, p=0.739</p> <p>2. Grp 1 vs. Grp 2, p&gt; 0.05</p> <p>3. Grp 1 vs. Grp 2, p&gt; 0.05</p> <p>4. Grp 1 vs. Grp 2, p&gt; 0.05</p>

	female, 3 male) (mean age 59.1 (9.3)), 7 in the NI group (3 female, 4 male) (mean age 58.9 (12.6))  <b>Setting:</b> rehabilitation hospital in Winnipeg, Manitoba	no intervention group (n=7)		
<b>Acupuncture</b>				
Kim 2016	50 participants mean age of 62.0±9.8 years, acupuncture (n=26), age 65.0±8.7, male / female 12/14, control (n=24), age 58.9±10.2, male / female 10/14. Mean duration of symptoms 33m  <b>Setting:</b> Hospital in Yangsan, South Korea	1) Acupuncture: 269 acupuncture sessions were administered during the study. 81% (n=21) of patients received at least 10 acupuncture sessions. Electrical acupuncture was applied at least once and bilaterally at back shu points (BL23, BL24, BL25 or BL26) or Jiaji points at L2– L5 spinal levels. Other frequently used points were BL57, BL60, GB39, GB34 and tender points located in the lower extremities (n=26)  2) Control: In total, 255 physical therapy sessions were provided to patients in the control group at their request. 92% (n=22) of patients received at least 10 physical therapy sessions (median 11, range 1–13). (n=24)	1) ODI 2) SF-36 bodily pain 3) SF-36 physical function 4) LBP bothersomeness 5) LBP intensity 6) Leg pain bothersomeness 7) Leg pain intensity 8) Self-reported pain-free walking distance (m)  <b>Follow-up:</b> 6 weeks, 3 months	<b>Between group MD, 95% CI</b> <b>ODI</b> 6 wk: -2.2 (-7.0 to 2.6) 3 mo: -2.5 (-8.9 to 3.8) <b>SF-36 BP</b> 6 wk: -8.6 (-18.6 to 1.3) 3 mo: 3.2 (-8.3 to 14.7) <b>SF-36 PF</b> 6 wk: 0.1 (-7.6 to 7.9) 3 mo: 1.3 (-8.3 to 10.9) <b>LBP bothersomeness</b> 6 wk: -0.6 (-11.4 to 10.1) 3 mo: -7.4 (-19.6 to 4.8) <b>LBP intensity</b> 6 wk: -5.1 (-15.5 to 5.3) 3 mo: -13.5 (-26.2 to -0.7) <b>Leg pain bothersomeness</b> 6 wk: -7.4 (-18.4 to 3.7) 3 mo: -9.2 (-21.6 to 3.2) <b>Leg pain intensity</b> 6 wk: -11.5 (-0.9 to -22.0) 3 mo: -12.6 (-24.6 to -0.6)

				None statistically significant
Qin 2020	<p>80 participants assigned with 70 completing the 8-week treatment course (38 in acu group and 32 in sham acu group). Mean age of 61.5±7.9 years with 34 males and 46 females. Duration of symptoms &lt;3mo =14 (17.5%), 3-12 mo = 1(1.3%), 1 to 5 y = 24 (30%), &gt;5 y =41 (51.3%)</p> <p><b>Setting:</b> 2 Clinical Sites - Department of Acupuncture and Neurology, Guang'anmen Hospital Department of Acupuncture and Neurology, Beijing Fengtai Hospital of Integrated Traditional and Western Medicine.</p>	<p>1) Acupuncture: Applied by acupuncturists with 5 years of Chinese medical university program and at least 2 year of clinical experience. Sterile disposable steel needles (Hwato Acupuncture, Suzhou, China; 0.30 £ 40 mm/0.30 £ 75 mm) were inserted through adhesive pads. Participants underwent 3 treatments weekly over 8 weeks, and each session persisted for 30 minutes. To maintain “De qi,” a sensation of numbness and soreness, acupuncture manipulation (twirling, lifting, and thrusting on needles) was performed every 10 minutes during the treatment.</p> <p>2) Sham acupuncture: Chosen acupoints, treatment duration, and frequency of sessions were the same as in the acupuncture group. Participants in the sham cohort were treated using a pragmatic placebo needle on the same acupoints, which is similar to the Streitberger needle design (Supplementary Materials). Acupuncturists pretended to manipulate the needle every 10 minutes, but “De qi” was not sought.</p>	<p>1) RMDQ 2) NRS back 3) NRS Leg 4) SSS Symptoms subscale 5) SSS physical function subscale 6) SSS satisfaction subscale 7) Self-paced walk test</p> <p><b>Follow-up:</b> 4 weeks, 8 weeks (end of treatment), 3 months, 6 months</p>	<p><b>RMDQ</b> 4 wk: -3.6 (-5.2 to -1.9): p&lt;0.001 8 wk: -2.6 (-3.7 to -1.4): p&lt;0.001 3 mo: -2.3 (-3.9 to -0.7): p=0.005 6 mo: -1.8 (-3.6 to -0.3): p=0.086</p> <p><b>NRS Back</b> 4 wk: -1.7 (-2.4 to -0.9): p&lt;0.001 8 wk: -2.3 (-3.0 to -1.5): p&lt;0.001 3 mo: -1.7 (-2.6 to -0.8): p&lt;0.001 6 mo: -1.2 (-2.1 to -0.3): p=0.007</p> <p><b>NRS Leg</b> 4 wk: -2.0 (-2.6 to -1.3): p&lt;0.001 8 wk: -2.9 (-2.6 to -1.3): p&lt;0.001 3 mo: -2.4 (-3.3 to -1.4): p&lt;0.001 6 mo: -2.1 (-3.0 to -1.2): p&lt;0.001</p> <p><b>SSS Symptoms Subscale</b> 4 wk: -0.6 (-0.8 to -0.4): p&lt;0.001 8 wk: -0.9 (-1.2 to -0.6): p&lt;0.001 3 mo: -0.9 (-1.2 to -0.6): p&lt;0.001 6 mo: -1.0 (-1.3 to 0.6): p&lt;0.001</p> <p><b>SSS Physical Function Subscale</b> 4 wk: -0.5 (-0.8 to -0.3): p&lt;0.001 8 wk: -0.8 (-1.1 to -0.5): p&lt;0.001 3 mo: -0.7 (-1.0 to -0.4): p&lt;0.001 6 mo: -0.7 (-1.1 to -0.4): p&lt;0.001</p> <p><b>Self-Paced Walk Test</b> 4 wk: p=0.648 8 wk: p=0.29 3 mo: p=0.30 6 mo: p=0.133</p> <p><b>Adverse events:</b> 3 participants in group 1 reported pain after needle insertion and 1 had a hematoma. 3 participants in group 2 reported back pain and 2 reported fatigue. All adverse events were reported as mild or moderate, and none required medical intervention.</p>



Epidural injections				
Cuckler 1985	<p>73 subjects in total, 37 with spinal stenosis, 36 with acute herniated nucleus pulposus, 37 males, 36 female, average age of 48.5 years in the experimental group and 49.5 years in the placebo group. Experimental group average 36.6 months in symptom duration, placebo group averaged 29.4 months.</p> <p><b>Setting:</b> Orthopaedic surgery department in the United States</p>	<p>1) Steroid group: 2ml of sterile water containing 80mg of methylprednisolone acetate combined with 5ml of 1% procaine was injected into the epidural space in the region between the 3rd and 4th lumbar vertebrae with the patient in the lateral decubitus position lying on the side of the painful limb (n=42), 20 with stenosis).</p> <p>2) Placebo group: 2ml of saline combined with 5ml of 1% procaine was injected into the epidural space in the region between the 3rd and 4th lumbar vertebrae with the patient in the lateral decubitus position lying on the side of the painful limb. (n=31, 17 with stenosis)</p> <p>All patients were advised to take mild analgesics (aspirin or acetaminophen) during the post-injection period. Second injection given if less than 50% improvement after 24 hours - considered treatment failure</p>	<p>1) Subjective percentage of improvement with 75% required to be considered a treatment improvement, if less than 50% after 24 hours was considered a treatment failure</p> <p>2) Re-injection rates</p> <p>3) Surgery rates</p> <p><b>Follow-up:</b> 24 hours, every 3 months up to 30 months, averaging 20.2 months in the steroid group and 21.5 months in the control group.</p>	<p><b>Patient Global Assessment (improved by at least 75%)</b></p> <p>24 hours: 33% (steroid) vs. 21% (saline) p&gt;0.05</p> <p>Long term: 33% (saline) vs. 14% (saline) p&gt;0.05</p>
Fukusaki 1988	<p>53 subjects, 38 males and 15 female. Group 1 averaged 70 years of age and 79 days of symptoms on average, group 2 averaged 69 years of age and</p>	<p>1) Epidural injection with 8 ml of saline, repeated twice in the first week (n=16)</p> <p>2) Epidural injection with 8 ml of 1% mepivacaine, repeated twice in the first week. (n=18)</p> <p>3) Epidural injection with a mixture of 8 ml of 1% mepivacaine and 40 mg</p>	<p>1) Walking distance which was graded according to distance (excellent, good, or poor)</p> <p><b>Follow-up:</b> 1 week, 1 month, 3</p>	<p><b>Walking distance</b></p> <p>Percent excellent effect = mean of &gt; 100m in walking distance</p> <p>1 week: 12.5 % (saline) vs. 55% (block) vs. 63.2% (block + steroid); block or block + steroid &gt; saline, p&lt; 0.05;</p> <p>1 mo: 6.3% (saline) vs. 16.7% (block) vs. 15.8% (block + steroid) p &gt; 0.05</p> <p>3 mo: 6.3 (saline) vs. 5.6% (block) vs. 5.3% (block +steroid) p&gt; 0.05</p> <p>No significant difference between block vs. block + steroid at</p>

	<p>an average of 82 days of symptoms, group 3 averaged 72 years of age and 94 days of symptoms on average</p> <p><b>Setting:</b> Anaesthesia department in Japan</p>	<p>of methylprednisone, repeated twice in the first week. (n=19)</p>	<p>months</p>	<p>all follow-up periods, <math>p&gt;0.05</math></p> <p><b>Adverse events:</b> no reported complications</p>
<p>Zahaar 1991</p>	<p>30 subjects, 37 male and 26 female. Steroid group averaged 46.5 years of age and 36.6 months of symptoms, control group averaged 49 years of age and 29.4 months of symptoms</p> <p><b>Setting:</b> Medical facility in Egypt</p>	<p>1) Steroid injection: 5ml of hydrocortisone acetate suspension, 2x2ml carbocaine, 4% Volume completed with sterile saline to 30ml (n=18)</p> <p>2) Control: 2x2ml of carbocaine, 4% injected into epidural space. Volume completed with sterile saline to 30ml. (n=12)</p>	<p>1) Subjective percentage of improvement where 75% or more was deemed successful and surgery after injection was considered a failure.</p> <p><b>Follow-up:</b> 24 hours, then every three months up to 36 mo averaging 20.2 mo in the steroid group and 21.5 mo control group.</p>	<p><b>Patient Global Assessment</b> (improved by at least 75%) 24 hours: 55% (steroid injection) vs. 50% (control) <math>p&gt;0.05</math> Up to 36 mo: 38% (steroid injection) group vs. 33.3% (control) <math>p&gt;0.05</math></p> <p><b>Failures (%)</b> (required surgery) Up to 36 mo: 61% (steroid injection) vs. 66.6% (control) <math>p&gt;0.05</math></p>
<p>Friedly 2014, 2017 Makris 2016</p>	<p>400 patients, 221 females and 179 males, 200 in the lidocaine group</p>	<p>1) Lidocaine + glucocorticoid (1-3 mL of 0.25-1% lidocaine followed by 1-3 mL triamcinolone (60-120mg), betamethasone (6-12mg),</p>	<p>1) RMDQ 2) NRS (Leg Pain)</p>	<p><b>Between group MD, 95% CI, p values</b> <b>RMDQ</b> 3 weeks: -1.8 (-2.8 to -0.9): <math>p&lt;0.001</math> 6 weeks: -1.0 (-2.1 to 0.1): <math>p=0.07</math></p>

	<p>with an average age of 68.1 years old and 200 glucocorticoid-lidocaine group with an average age of 68 years old, LSS by CT or MRI. 26% patients symptoms greater than 5 years.</p> <p><b>Setting:</b> 16 medical centers across the United States</p>	<p>dexamethasone (8-10mg) or methylprednisone (60-120mg)) (n=200)</p> <p>2) Lidocaine group (0.25-1% lidocaine alone) (n=200)</p> <p>Physician option for intralaminar and/or transformaminar techniques</p>	<p><b>Follow-up:</b> 3, 6, and 12 weeks, 6 and 12 months</p> <p>Makris 2016 subgroup</p> <p>1) RMDQ using SIP Weights</p> <p>2) RMDQ patient-prioritized (LESSER)</p> <p><b>Follow-up:</b> 3 and 6 weeks</p>	<p>12 wk: 0.1 (-1.0 to 1.3): p=0.84 6 mo -0.00 (-1.1 to 1.1): p=0.99 12 mo: -0.4 (-1.6 to 0.9): p=0.55</p> <p><b>NRS (Leg pain)</b></p> <p>3 weeks: -0.6 (-1.2 to -0.1): p=0.02 6 weeks: -0. (=0.8 to 0.4): p=0.48 12 wk: 0.1 (-0.5 to 0.7): p=0.70 6 mo: -0.2 (-0.8 to 0.4): p=0.47 12 mo: 0.1 (-0.5 to 0.7): P=0.75</p> <p>Subgroup Analysis</p> <p><b>RMDQ using SIP weight</b></p> <p>3 wks: -1.9 (-2.9 to -0.7): p&lt;0.001 6 wks: -1.1 (-2.2 to -0.1): p=0.04</p> <p><b>RMDQ patient prioritized (LESSER)</b></p> <p>3 wks: -1.8 (-2.8 to -0.8): p&lt;0.001 6 wks: -1.0 (-2.0 to 0.1): p=0.08</p> <p><b>Adverse events:</b> A total 21.5% of patients in group 1 and 15.5% in group 2 reported one or more adverse events (p=0.08) that included headaches, fever, infection, dizziness, cardiovascular/lung problems, leg swelling and dural puncture.</p>
Song 2016	<p>29 subjects, 14 males and 15 women with an average age of 58.3 and 61.7 between groups, history of intermittent claudication and lower limb radicular pain or paresthesia</p>	<p>1) Lidocaine spinal injection, 40 mg triamcinolone mixed with 10 mL 0.5% lidocaine was used under the guide of fluoroscopy (n=15)</p> <p>2) Saline spinal injection using same volume (n=14)</p>	<p>1) VAS</p> <p>2) FRI</p> <p><b>Follow-up:</b> 1 and 3 months</p>	<p>No significant difference between groups.</p> <p><b>VAS</b></p> <p>1-month p= 0.696, 3 months p= 0.891</p> <p><b>FRI</b></p> <p>1-month p=0.983, 3 months p=0.743</p>

	<b>Setting:</b> Rehabilitation clinic in Korea			
Milburn 2014	<p>57 patients met inclusion criteria, agreed to participate, and were enrolled. 20 patients were male; 37 were female. Mean patient age was 65.3 years (range, 32-88 years). Average duration of symptomatology (pain and/or disability) was 42 months. The mean degree of canal narrowing at the most stenotic level was 6.1 mm (range, 2.5-9.1 mm). The most common maximally stenotic intervertebral level was L4-L5</p>	<p>Fluoroscopically guided lumbar ILES performed either at:</p> <ol style="list-style-type: none"> <li>1) The level of maximal stenosis (n=30)</li> <li>2) Two intervertebral levels cephalad, corresponding to a less stenotic level (n=27)</li> </ol> <p>Injection was performed with a 20-gauge Tuohy needle using a loss of resistance technique. The injectate consisted of 2 mL of 40 mg/mL methylprednisolone (Pfizer), 2 mL of bupivacaine 0.25% (Hospira), and 2 mL of normal saline for a total injectate volume of 6 mL.</p>	<ol style="list-style-type: none"> <li>1) NRS - Pain with Ambulation</li> <li>2) RMDQ</li> </ol> <p><b>Follow-up:</b> 1, 4 and 12 weeks</p>	<p><b>All between group comparisons</b></p> <p><b>NRS (pain with ambulation)</b></p> <p>1 wk: Grp 1 lower pain compared to Grp 2, p=0.045</p> <p>4 wk: Grp 1 lower pain compared to Grp 2, p=0.049</p> <p>12 wk: Grp 1 lower pain compared to Grp 2, p=0.08</p> <p><b>RMDQ</b></p> <p>1 wk: Grp 1 lower compared to Grp 2, p=0.001</p> <p>4 wk: Grp 1 lower compared to Grp 2, p=0.009</p> <p>12 wk: Grp 1 lower compared to Grp 2, p=0.003</p>

	(n/42) followed by L3-L4 (n/41) and L5-S1 (n/4).			
	<b>Setting:</b> Clinic in New Orleans, Louisiana			
Brown 2012	38 patients, 21 males and 17 females, 21 in mild group with an average age of 74.2 years and 17 in ESI group with an average age of 78.7 years, symptomatic LSS patients with painful lower limb neurogenic claudication, able to walk at least 10 feet unaided, (ODI) score > 20  <b>Setting:</b> Pain management clinic in Florida	1) Epidural steroid (80 mg triamcinolone acetate) (n=17) 2) Mild lumbar decompression (n=21)	1) VAS 2) ODI 3) ZCQ 4) Patient Satisfaction (0-10)  <b>Follow-up:</b> 6 and 12 weeks	<b>VAS</b> 6 and 12 weeks P=0.54 <b>ODI</b> p=0.86 <b>ZCQ</b> p>0.05 <b>Patient satisfaction</b> p>0.05
Hammerich 2019	54 patients total, age 67.2 ± 9.7, 27 male, 27 female, 31 in ESI group, 23 in ESI plus PT. Mean duration of	1) ESI (n=31) 2) ESI + PT (n=23)  ESI: 1.5 mL of steroid at each site injected with maximal involvement using transforaminal approach.	1) ODI 2) NRS current 3) SF-36 emotional role 4) SF-36 emotional well-being	<b>Between group MD, 95% CI, p values</b> <b>ODI</b> 10 wks: -1.08 (-8.10 to 5.94) p=0.80 6 mo: -4.70 (-11.72 to 2.32) p=0.27 12 mo: -2.72 (-9.74 to 4.30) p=0.52 <b>NRS</b> 10 wks: -1.68 (-3.08 to -0.29) p=0.07

	<p>symptoms 14 m</p> <p><b>Setting:</b> Clinics in Colorado, Texas, South Carolina and New Hampshire</p>	<p>PT: 8-10 sessions PT manual therapy and exercise. Walking program and/or stationary bike, stretching and strengthening exercises.</p>	<p>5) SF-36 general health perception</p> <p><b>Follow-up:</b> 10 weeks, 6 and 12 months</p>	<p>6 mo: -1.99 (-3.38 to -0.60) p=0.04 12 mo: -2.44 (-3.80 to -1.08) p=0.00</p> <p><b>SF-36 Emotional role</b> 10 wks: -28.53 (-49.05 to -8.01) p=0.03 6 mo: -11.25 (-31.77 to 9.27) p=0.39 12 mo: -10.67 (-31.19 to 9.85) p=0.41</p> <p><b>SF-36 Emotional well-being</b> 10 wks: -11.26 (-19.52 to -2.99) p=0.02 6 mo: 2.69 (-5.57 to 10.95) p=0.59 12 mo: -5.76 (-14.02 to 2.50) p=0.24</p> <p><b>SF-36 General Health Perception</b> 10 wks: -8.99 (-17.20 to -0.78) p=0.05 6 mo: -5.56 (-13.77 to 2.65) p=0.23 12 mo: -5.10 (-13.31 to 3.11) p=0.27</p>
Sencan 2020	<p>67 patients. The median age 62.5 years with 18 males and 49 females. Median duration of symptoms was 29 and 24 months in the ILESI and bilateral TFESI groups, respectively</p> <p><b>Setting:</b> University department Pain Medicine, Istanbul Turkey</p>	<p>1) Interlaminar: ILESI, fluoroscopy guided with 1 to 2 mL contrast dye with mixture of 80 mg methylprednisolone acetate, 2 mL saline solution, and 2 mL (0.5%) bupivacaine solution</p> <p>2) Transforaminal: TFESI, fluoroscopy guided with 1 to 2 mL contrast dye with mixture of 80 mg methylprednisolone acetate, 2 mL saline solution, and 2 mL (0.5%) bupivacaine solution</p>	<p>1) NPS 2) ODI 3) Beck depression scale 4) Walk distance</p> <p><b>Follow-up:</b> after treatment, 3 weeks and 3 months</p>	<p><b>Between Group Median Differences (data not provided), p values</b> <b>NPS</b> after treatment: p=0.14 3 wks: p=0.28 3 mo: p=0.047</p> <p><b>ODI</b> 3 wks: p=0.93 3 mo: p=0.65</p> <p><b>Beck Depression Scale</b> 3wks: p=0.048 3 mo: p=0.03</p> <p>Walking Distance 3 wks: p=0.23 3 mo: p= 0.048</p>
Wei 2020	<p>90 patients. Mean age about 65 years, 45 females, 45</p>	<p>1) Epidural injection with 2.0mL of lidocaine and 10 mg of TNF-<math>\alpha</math> inhibitor (etanercept) on the affected spinal nerves.</p>	<p>1) VAS (leg) 2) ODI</p> <p><b>Follow-up:</b> after</p>	<p><b>Between Group Mean Differences (data not provided), p values</b> <b>Grp 1 vs Grp 2</b> <b>VAS</b></p>



	<p>males, mean duration of symptoms about 2.8 months</p> <p><b>Setting:</b> University Hospital Jiangsu China</p>	<p>2) Epidural administration with 2mL of lidocaine mixed with 2mL of steroid (diprosan)</p> <p>3) Epidural injection 4.0mL of lidocaine only.</p>	<p>treatment, 1,3, 6 months</p>	<p>after treatment, 1, 3 and 6 mo, Grp 1 greater reduction, <math>p&lt;0.05</math></p> <p><b>ODI</b></p> <p>1, 3 and 6 mo, Grp 1 greater reduction, <math>p&lt;0.05</math></p> <p><b>Grp 1 vs Grp 3</b></p> <p><b>VAS</b></p> <p>after treatment, 1, 3 and 6 mo, Grp 1 greater reduction, <math>p&lt;0.05</math></p> <p><b>ODI</b></p> <p>1, 3 and 6 mo, Grp 1 greater reduction, <math>p&lt;0.05</math></p> <p><b>Grp 2 vs Grp 3</b></p> <p><b>VAS</b></p> <p>after treatment, 1, 3 and 6 mo, no significant difference, <math>p&gt;0.05</math></p> <p><b>ODI</b></p> <p>1, 3 and 6 mo, no significant difference, <math>p&gt;0.05</math></p>
Karm 2018	<p>44 patients total, 20 in the RACZ group (age 66.1 <math>\pm</math> 12.2, male 9 (45.0%), and 24 in the ZiNeu group (Age 65.5 <math>\pm</math> 6.4 18 females, 26 males.</p> <p><b>Setting:</b> Single-center, academic, outpatient interventional pain management clinic in Korea</p>	<p>1) PEA Using a Balloon-less Catheter (Racz) (n = 20)</p> <p>2) Percutaneous Epidural Decompression and Adhesiolysis Using an Inflatable Balloon Catheter (ZiNeu) (n = 24)</p>	<p>1) NRS (back pain)</p> <p>2) NRS (leg pain)</p> <p>3) ODI</p> <p><b>Follow-up:</b> 1, 3 and 6 months</p>	<p><b>Between group MD, 95% CI, p values</b></p> <p><b>NRS-11 (Back pain)</b></p> <p>1 mo: -0.38 (-1.81 to 1.06): <math>p=0.61</math></p> <p>3 mo: -1.13 (-2.63 to 0.38): <math>p=0.14</math></p> <p>6 mo: -2.02 (-3.58 to 0.45): <math>p=0.01</math></p> <p><b>NRS-11 (Leg pain)</b></p> <p>1 mo: 0.73 (-0.40 to 1.85): <math>p=0.21</math></p> <p>3 mo: -0.69 (-1.89 to 0.52): <math>p=0.26</math></p> <p>6 mo: -1.88 (-3.15 to 0.61): <math>p=0.00</math></p> <p><b>ODI (%)</b></p> <p>1 mo: -6.13 (-13.88 to 1.61): <math>p=0.12</math></p> <p>3 mo: -6.63 (-14.75 to 1.48): <math>p=0.11</math></p> <p>6 mo: -13.74 (-22.18 to 5.30): <math>p=0.00</math></p> <p><b>Adverse events:</b> Minor and transient adverse events were reported equally in both groups (no data provided), mostly pain and paresthesia at the injection site.</p>
<b>Surgery</b>				
Zucherman 2004, 2005, 2006	<p>191 subjects, 57% male and 43% female in the X STOP group. 52% male</p>	<p>1) X STOP Interspinous Process Decompression System (n=100)</p> <p>2) Non-operative treatment: Subjects received an epidural steroid injection</p>	<p>1) SF-36</p> <p>2) ZCQ</p> <p>3) Worker's compensation claims</p>	<p><b>Patient global assessment (Good result)</b></p> <p>2 yrs: 73.1% (surgery) vs. 35.9% (control) (<math>P&lt;0.001</math>)</p> <p><b>Symptoms Severity score</b></p>

	<p>and 48% female in the non-operative group. Average age of 70 years in the X STOP group and 69.1 years in the non-operative group. Average of 3.5 year symptom duration in the X STOP group and 4.7 years in the non-operative group.</p> <p><b>Setting:</b> Spine center in the United States</p>	<p>on enrolment and were eligible for additional injections as needed, as well as NSAIDS, analgesic agents, and physical therapy. Physical therapy consisted of education on back care and modalities such as ice packs, heat packs, massage, stabilization exercises, and pool therapy. Braces such as abdominal binders and corsets were permitted, but body jackets and chair back braces were not. (n=91)</p>	<p>4) ODI 5) Radiographic changes</p> <p><b>Follow-up:</b> Surgery: 7 (2 yr) Control: 19 (2 yr)</p>	<p>Surgery better at 6 w, 6 mo, 1 and 2 yr (graphs) (P&lt;0.001) 2 yrs: MPC 45.4% (surgery) vs. 7.4% (control) (P &lt; 0.001) <b>“Clinically relevant improvement (patients)”:</b> 2 yrs: 60.2% (surgery) vs. 18.5% (control) (P&lt; 0.001) <b>Symptoms Severity score††</b> Surgery better at 6 w, 6 mo, 1 and 2 yr (graphs) (P&lt;0.001) 2 yrs: MPC 44.3% (surgery) vs. -0.4% (control) (P &lt; 0.001) <b>“Clinically relevant improvement (as measured by patients)”:</b> 2 yrs: 57% (surgery) vs. 14.8% (control) (P &lt; 0.001) <b>ZCQ (global success)</b> 6 mo: 52% (surgery) vs. 9% (control) (P value not reported) 1 yr: 59% vs 12% (P value not reported) 2 yrs: 48.4% (surgery) vs. 4.9% (control) (P &lt; 0.001) <b>Quality of life (SF-36)</b> At all post treatment time points (6 w, 6 mo, 1 yr, 2 yr), the mean domain scores documented in the X STOP group were significantly greater than those in the non operative group, with the exception of the mean General Health, Role Emotional, and Mental Component <i>Summary scores at 2 years</i></p> <p><b>Adverse events:</b> No complications were reported in group 2. In group 1, complications were reported in 11% of subjects including spinous process fracture, coronary ischemia, respiratory distress, hematoma, and 1 death (pulmonary edema)</p>
Weinstein 2007, 2009, Abdu 2018	<p>Subjects with image-confirmed degenerative spondylolisthesis: 304 subjects in the RCT, 303 in the observational cohort, 31% male in the surgical group, 33% male in the surgical group. Average</p>	<p>1) Assigned to surgery (standard laminectomy with or without fusion) (n=159) 2) Assigned to non-surgical treatment: Usual non-operative care (n=145)</p>	<p>1) SF-36 bodily pain 2) SF-36 bodily function 3) low back pain bothersomeness scale 4) Leg pain bothersomeness scale 5) ODI 6) Subjective self-</p>	<p><b>All between group comparisons using Intention-to-Treat analysis</b> <b>SF-36 Bodily Pain, DMC, 95% CI</b> 2 yrs: 1.5 (-4.2 to 7.3) 4 yrs: -2 (-8.6 to 4.6) 8 yrs: p=0.85 <b>SF-36 Bodily Function, DMC, 95% CI</b> 2 yrs: 1.9 (-3.7 to 7.5) 4 yrs: -3.1 (-9.2 to 3.0) 8 yrs: p=0.31 <b>Disability (ODI), DMC, 95% CI</b> 2 yrs: 2.2 (-2.3 to 6.8)</p>

	age of 64.7 years in the surgical group and 68.2 years in the non-surgical group. Subjects had symptoms for at least 12 weeks  <b>Setting:</b> multi-centred orthopaedic departments in the United States		reported improvement, satisfaction with current symptoms and care 7) Stenosis bothersomeness index  <b>Follow-up:</b> 6 weeks, 3 and 6 months, 1, 2, 4 and 8 years	4 yrs: 4.1 (-0.8 to 9.1) 8 yrs: p=0.039  Other outcomes (patient's satisfaction; Stenosis Bothersomeness Index, Leg Pain Bothersomeness Scale; and Low Back Pain Bothersomeness Scale) were not provided separately for the randomized cohort. <b>Adverse events:</b> group 1 reported 14% intraoperative complication mostly and dural tears and 19% postsurgical complications including 1 death, 11% required additional surgeries at 2 years,
Amundsen 2000	100 subjects, 54 male, 46 female, median age of 59 (males were 1.5 years higher than females). Median back pain duration was 14 years, median duration of sciatica was 2 years.  <b>Setting:</b> Neurology department in a hospital in Norway	1) Surgery: Partial or total laminectomy, medial facetectomy, discectomy, and/or removal of osteophytes from the vertebral margins or facet joints. No fusions. (n=13)  2) Conservative therapy: Lumbar orthosis use for 1 month worn during the day for all activities plus instruction and back school." (n=18)	1) VAS 2) Verbal Rating Scale 3) Subjective change (better, worse, or unchanged) 4) Work status 5) Subjective rating from evaluating physician and study team (Excellent, Fair, Unchanged, Worse)  <b>Follow-up:</b> 6 months, 1, 4 and 10 years	<b>Patient global assessment (Good result)</b> 1 yr: RR 2.07 (0.98 to 4.38) 4 yrs: RR 1.94 (1.14 to 3.31) 10 yrs: RR 3.18 (0.97 to 10.41) <b>Pain (none or mild)</b> 1 yr: NR 4 yrs: RR 3.33 (0.77 to 14.33) 10 yrs: RR 1.59 (0.55 to 4.55)  Other outcomes (claudication or walking distance; level of daily activity; and neurologic deficits) were not reported separately for the randomized cohort.
Malmivaara 2007	94 subjects, 22% of surgical	1) Segmental decompressive surgery with facetectomy (n=50)	1) 11 point numerical pain	<b>All between group comparisons</b> <b>Leg pain, MD, 95% CI</b>

	<p>subjects were male, 45% of non-operative subjects were male. Nonoperative group had average age of 62.9 years, surgical group had average age of 63.9 years. Surgical group averaged 14 years since onset of symptoms, nonsurgical group average 16 years since onset of symptoms. Minimum of 6 months of symptoms for study inclusion.</p> <p><b>Setting:</b> Research Center in Finland</p>	<p>2) Non-operative treatment: NSAIDS when indicated and seen one to three times by a physiotherapist, in addition to the standard visit at each follow-up. The physiotherapist gave all patients educational brochure. The patients were encouraged to use their back in a normal way. Pain-relieving body postures were taught as well as basic ergonomics related to lifting and carrying. Individually structured programs included trunk muscle endurance and stretching-type exercises. Additional individual physiotherapy consisting of passive treatment methods (such as ultrasound and transcutaneous nerve stimulation). (n=44)</p> <p>The patients in the surgical group also received the brochure and the instructions described above.</p>	<p>rating scale for back and leg pain</p> <p>2) Walking ability (distance without a break) also via treadmill test</p> <p>3) General health status on a 5 point scale (very good, quite good, average, quite poor or very poor.</p> <p>4) ODI</p> <p>5) Ability to complete certain activities of daily</p> <p>6) living without difficulty, some difficulty, marked difficulties or not at all</p> <p>7) Radiographic examination</p> <p><b>Follow-up:</b> 6 months, 1 and 2 years</p>	<p>1 yr: 1.69 (0.41 to 2.96) 2 yr: 1.51(0.25 to 2.77) <b>Back pain, MD, 95% CI</b> 1 yr: 2.33 (1.12 to 3.55) 2 yrs: 2.13(0.98 to 3.28) <b>Disability (ODI), MD, 95% CI</b> 1yr: 11.3 (4.3to 18.8) 2 yrs: 7.8 (0.8 to14.9) <b>&gt; 10 points reduction (ODI): RR, 95% CI</b> 1 yr: 2.16 (1.31to 3.57) 2 yrs: 1.36 (0.88 to 2.10)</p> <p><b>Walking disability (walking distance &lt;1.250 m), RR, 95% CI</b> 1 yr: 0.93 (0.61 to 2.03) 2 yrs: 1.08 (0.70 to 2.42) <b>Walking disability (walking distance &lt;400 m), RR, 95% CI</b> 1 yr: 0.91 (0.51 to 4.24) 2 yrs: 1.18 (0.67 to 4.72)</p>
--	--	--	---	--

Weinstein 2008, 2010, Lurie 2015	<p>289 in the RCT, 365 in the observational cohort. 62% male in the surgical groups, 59% male in the non-surgical groups. Average age of 63.8 in the surgical group, 66.1 in the non-surgical group. 60% in the surgical group and 55% in the non-surgical group had symptoms for over 6 months.</p> <p><b>Setting:</b> multi-centred-orthopaedic departments in the United States.</p>	<p>1) Assigned to surgery: Standard laminectomy with or without fusion (n=138)</p> <p>2) Assigned to non-surgical treatment: Usual non-operative care - recommended to include at least active physical therapy, education or counseling with home exercise instruction, and the administration of NSAIDs, if tolerated (n=151)</p>	<p>1) SF-36 bodily pain</p> <p>2) SF-36 bodily function</p> <p>3) Low back pain bothersomeness scale</p> <p>4) Leg pain bothersomeness scale</p> <p>5) ODI</p> <p>6) Subjective self-reported improvement, satisfaction with current symptoms and care,</p> <p>7) Stenosis bothersomeness index</p> <p><b>Follow-up:</b> 6 weeks, 3 and 6 months, 1, 2, 4, 8 years</p>	<p><b>All between group comparisons using Intention-to-Treat Analysis</b></p> <p><b>SF-36 Bodily Pain, DMC, 95% CI</b> 2 yrs: 7.8 (1.5 to 14.1) 4 yrs: 0.3 (-6.4 to 7) 8 yrs: p=0.25</p> <p><b>SF-36 Bodily Function, DMC, 95% CI</b> 2 yrs: 0.1 (-6.4 to 6.5) 4 yrs: -3.2 (-9.9 to 3.6) 8 yrs: p=0.89</p> <p><b>Disability (ODI), DMC, 95% CI</b> 2 yrs: -3.5 (-8.7 to 1.7) 4 yrs: 0.2 (-5.2 to 5.7) 8 yrs: p=0.87</p> <p>Other outcomes (patient's satisfaction; Stenosis Bothersomeness Index, Leg Pain Bothersomeness Scale; and Low Back Pain Bothersomeness Scale) were not provided separately for the randomized cohort.</p> <p><b>Adverse events:</b> In group 1, 10% of patients required transfusions intraoperatively and 5% postoperatively. The most common surgical complication was dural tear, in 9% of patients. At 2 years, reoperation had occurred in 8% of subjects.</p>
Delitto 2015	<p>169 patients, 88 males and 81 females, 87 surgical group with an average age of 66.6 years old and 82 PT group with an average age of 69.8 years old, LSS by computed</p>	<p>1) Surgical decompressive laminectomies, partial facet resection, and neuroforaminotomies (n=87)</p> <p>2) PT program: lumbar flexion exercises, exercises and education (n=82)</p>	<p>1) SF-36 physical function</p> <p><b>Follow-up:</b> 2 years</p>	<p><b>2 years -SF-36 Physical Function, MD and 95% CI</b> 0.9 (7.9 to 9.6)</p> <p><b>Adverse events:</b> 9 out of 82 participants in group 2 reported adverse events consisting of worsening of symptoms whereas 33 out of 87 participants in group 1 reported surgery related complications, mainly attributable to reoperation, delay in wound healing and surgical site infection.</p>

	tomography - criteria of Wiesel and colleagues (18) or magnetic resonance imaging - criteria of Boden and colleagues (2)  <b>Setting:</b> Neurologic and orthopedic surgery departments and physical therapy clinics in western Pennsylvania			
--	--	--	--	--

ADLs = Activities of Daily Living, AUC = Area under the pain-intensity curve, BTX = Botox, CI = Confidence Interval, DMC = Difference in mean change from baseline, ESI = Epidural Steroid Injection, FRI = Functional Rate Index, GRP = Group, HADS = Hospital Anxiety and Depression Scale, IU = International Units, JOABPEQ = Japanese orthopaedic association back pain evaluation questionnaire, LBOS = Low Back Outcome Score, LBP = Low Back Pain, m = Meters, MCS = Mental Component Score, MD = Mean Difference, mm = Millimeters, Mo = Months, MPC = Mean Percent Change, NRS = Numerical Pain Rating Scale, NR = Not Reported, ODI = Oswestry Disability Index, OR = Odds Ratio, PASS-20 = Pain Anxiety Symptoms Scale, PCS = Physical Component Score, RCT = Randomized Controlled Trial, RMDI = Roland Morris Disability Index, ROM = Range of Motion, RR = Relative Risk, SBI = Stenosis Bothersomeness Index, SPWT = Self-Paced Walking Test, SSS = Spinal Stenosis Questionnaire, TSK-11 = Tampa Scale-11, VAS = Visual Analogue Scale, WMD = Weighted Mean Difference, ZCQ = Zurich Claudication Questionnaire