

BMJ Open Protocol for a systematic review on routine use of antibiotics for infants less than 6 months of age with growth failure/faltering

Aamer Imdad ¹, Fanny F Chen,² Melissa François,² Emily Tanner-Smith ³, Abigail Smith,⁴ Olivia J Tsistinas,⁴ Jai K Das ⁵, Zulfiqar Ahmed Bhutta^{6,7}

To cite: Imdad A, Chen FF, François M, *et al.* Protocol for a systematic review on routine use of antibiotics for infants less than 6 months of age with growth failure/faltering. *BMJ Open* 2022;**12**:e057241. doi:10.1136/bmjopen-2021-057241

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-057241>).

Received 09 September 2021
Accepted 13 June 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Aamer Imdad;
imdada@upstate.edu

ABSTRACT

Introduction Antibiotics have been used as an adjunct in treating children with severe acute malnutrition 6–59 months of age; however, the data for infants less than 6 months are scarce. The WHO recently started guideline development for preventing and treating wasting, including growth failure/faltering in infants less than 6 months. This systematic review commissioned by WHO aims to synthesise evidence from current literature on the effectiveness of antibiotics for infants less than 6 months of age with growth failure/faltering.

Methods and analysis We will conduct a systematic review and meta-analysis for studies that assessed the effect of antibiotics in the treatment of infants with growth faltering. We will search multiple electronic databases. We will include randomised control trials and non-randomised studies with a control arm. The study population is infants less than 6 months of age with growth failure. The intervention group will be infants who received no antibiotics or antibiotics other than recommended in 2013 guidelines by WHO to treat severe acute malnutrition in children. The comparison group will be infants who received antibiotics according to the 2013 guideline by WHO. We will consider the following outcomes: mortality, clinical deterioration, antimicrobial resistance, recovery from comorbidity, adverse events, markers of intestinal inflammation, markers of systemic inflammation, hospital-acquired infections, non-response. We will use the meta-analysis to pool the studies where applicable. We will use the Grading of Recommendations Assessment, Development, and Evaluation approach to reporting the overall evidence quality for an outcome.

Ethics and dissemination This is a systematic review and will not involve contact with a human subject. The findings of this review will be published in a peer-review journal and will guide the WHO's recommendation for the use of antibiotics in infants less than 6 months of age with growth failure.

PROSPERO registration number CRD42021277073.

INTRODUCTION

The WHO and the United Nations Children's Fund (UNICEF) estimate that nearly 14 million children suffer from severe acute malnutrition (SAM) worldwide.¹ Infants less

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ We will search several electronic databases and include randomised and non-randomised studies.
- ⇒ Two authors will screen the titles and extract the data from included studies duplication. We will assess the risk of bias for each outcome from each study and use the Grading of Recommendation, Assessment, Development and Evaluation criteria to assess the overall quality of evidence.
- ⇒ We will conduct the meta-analysis if the data are available for more than one study for an outcome and there is clinical and methodological homogeneity in the included studies.
- ⇒ All the included studies may not report the data for all the outcomes mentioned in this review, which will be a limitation of the study.

than 6 months of age are particularly vulnerable to the effects of inadequate nutrition. Higher mortality rates secondary to growth failure are seen in this age group compared with older infants and children.^{2–3} Despite excess mortality risk and increasing prevalence of wasting in this population, limited studies exist to guide the management of young infants with growth failure and faltering.^{2–6} Malnutrition not only increases the risk of severe infections in children and triples the risk of mortality from pneumonia, measles or diarrhoea but also detrimentally affects their neurodevelopment.^{7–9} Therefore, the current practice for children 6 months to 5 years of age with SAM is to prescribe routine antibiotics, such as 5–7 days of amoxicillin, when they are inpatients.⁹ However, current recommendations state that the same general medical care should be used for infants with SAM who are less than 6 months of age as infants above 6 months of age, even though there is limited evidence to support this recommendation for infants less than 6 months of age.²

Furthermore, even though antibiotics have been shown to be effective in children 6–59 months of age with SAM, this practice in infants has the potential to harm due to recently identified risks of antibiotic use in infancy, including the diminishment of infant gut microbiome,¹⁰ future development of obesity, allergic disorders¹¹ and autoimmune disorders.¹² The urgency in more targeted management guidelines is further underscored by the physiological differences in renal and gastrointestinal function in infants compared with older children.¹² The WHO recently started the process of guideline development for the prevention and treatment of wasting in children, including growth failure/faltering in infants less than 6 months. We aim to systematically review and synthesise evidence from current literature on the effectiveness of antibiotics for infants less than 6 months of age with growth failure/faltering.

REVIEW QUESTION

In infants <6 months with growth failure/faltering, what are the effects of no routine antibiotics or different approaches (eg, types of antibiotics, doses, etc) compared with routine antibiotics following treatment protocols in the 2013 WHO guidelines on the outcomes of interest?

METHODS

This systematic review will be conducted according to methods described in Cochrane Handbook¹³ and reported using Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines 2020.¹⁴ We registered the protocol on the PROSPERO website.

Inclusion criteria

We will include randomised and non-randomised controlled trials (RCTs). We will consider both individual and cluster randomised trials. We will also consider non-randomised trials and cohort studies with a controlled arm. Table 1 gives the definitions of the eligible study designs. We will exclude case-control studies, case reports, case series and commentaries.

Population

The population of interest is infants under 6 months of age with growth failure/faltering. As there is no standard definition of growth failure/faltering, we will consider author definitions. We will include studies irrespective if they are done in the community setting or the hospital setting. We will consider studies that included infants infected with HIV. We will include studies with infants born low birth weight or preterm; however, we will exclude studies on infants admitted to the neonatal intensive care units. We will exclude studies that included infants with congenital anomalies only. If a study includes infants less than 6 months and greater than 6 months of age, we will consider this study if the disaggregated data are available for infants less than 6 months. If the disaggregated data are not available from such a study, we will consider the study if the average age of infants is at or below 6 months of age.

Intervention

We will include all antibiotic treatments given systematically, such as amoxicillin, augmentin, cephalosporins and macrolides. We will include studies irrespective of dosage, frequency, duration or route of administration; however, topical application of antibiotics will not be

Table 1 Suggested study designs for inclusion in the review

Suggested terms	Definition	Notes
Randomised controlled trial (RCT) OR randomised trial	An experimental study in which people are allocated to different interventions using methods that are random.	We will consider both individual and cluster randomised trials. We also consider the factorial design trials
Non-randomised controlled trial (NRCT) OR non-randomised trial	An experimental study in which people are allocated to different interventions using methods that are not random.	We will use consistent terminology and avoid using the term quasi-experimental as it might have different meanings in different settings. We will use the ROBINS-1 for risk of bias assessment for this study design
Controlled before-after study (CBA)	A study in which observations are made before and after the implementation of an intervention, both in a group that receives the intervention and in a control group that does not.	We will require two minimum criteria for the inclusion of CBAs. <ul style="list-style-type: none"> ► Data Collection: We will include CBAs if the data for the intervention and control groups were collected prospectively in the same time frame ► Choice of Control: We will include CBAs that include a control at a second site to avoid contamination of the intervention to the control group if the settings and populations are the same for the intervention and control groups.

This table is modified from: <https://epoc.cochrane.org/sites/epoc.cochrane.org/files/public/uploads/EPOCStudyDesignsAbout.pdf>

considered. We will consider studies if the antibiotics were given empirically at the diagnosis of growth failure or faltering. We will also consider studies if the antibiotics were given in response to a suspected infection in infants with wasting. We will exclude studies where antibiotics were given for the prevention of wasting in otherwise healthy infants. We will also exclude studies where antibiotics were given for other reasons, such as suspected serious bacterial infections in otherwise healthy infants. We will allow cointervention such as nutritional supplementation for nutritional rehabilitation only if similar in both groups.

Comparison

The comparison groups are routine antibiotics following treatment protocols detailed in the 2013 WHO guideline.²

Outcomes

- Mortality (dichotomous outcome).
- Clinical deterioration (dichotomous outcome, defined by the development of any danger signs (obstructed breathing, respiratory distress, cyanosis, shock, severe anaemia, convulsion, severe dehydration, profuse watery diarrhoea, vomiting and/or impaired consciousness)).
- Antimicrobial resistance (dichotomous outcome, as defined by authors).
- Recovery from wasting (dichotomous outcome, as defined by authors).
- Recovery from comorbidity, pneumonia (dichotomous outcome).
- Recovery from comorbidity, diarrhoea (dichotomous outcome).
- Recovery from any comorbidity (dichotomous outcome).

- Adverse events—incidence of diarrhoea (dichotomous outcome, >3 loose stools/24 hours).
- Markers of intestinal inflammation—faecal calprotectin (continuous outcome).
- Markers of systemic inflammation—serum C reactive protein (continuous outcome).
- Hospital-acquired infections (dichotomous outcome).
- Non-response (eg, not achieving recovery within 4 months of initiating treatment) (dichotomous outcome).

We will consider all the outcomes at 1, 3 and 6 months and the longest follow-up. All the primary analyses will be done at the longest follow-up.

Balance of benefits and harms

We will stratify the outcomes by benefit and harm and provide a narrative summary of the balance of benefits and harms.

Literature search

We will conduct systematic electronic queries using key terms in multiple databases, including MEDLINE via PubMed, EMBASE, Web of Science, CINAHL, Scopus, LILACS, WHO Global Index Medicus and BIOSIS Previews. There will be no search restrictions on outcomes, publication year, publication status or publication language. The search strategies for different databases are available in online supplemental appendix 1. The references of formerly published reviews and recently published studies will be examined for potential inclusion. We will also use The Cochrane Central Register for Controlled Trials and ISRCTN registry to identify studies currently underway. We will also search the websites of pertinent international agencies such as the WHO (including WHO's Reproductive Health Library, electronic Library of Evidence of Nutrition Actions and Global database on the Implementation of Nutrition Action), UNICEF, Global Alliance for Improved Nutrition, International Food Policy Research Institute, International Initiative for Impact Evaluation, Nutrition International, World Bank, USAID and affiliates (eg, FANTA, SPRING) and the World Food Programme. We will search the abstracts of the major conference, such as annual paediatric academic society meetings. Finally, we will use the citation tracking function of the included studies in PubMed to look for any eligible studies.

Data extraction and synthesis

Selection of studies and data extraction

We will use a three-phased approach to screen studies for eligibility. In the first phase, two authors will independently screen study titles and abstracts yielded from search results to identify potentially eligible ones. Studies selected during this initial phase will then go through a full-text review, which constitutes the second phase. In the final phase, studies that are determined to be eligible through the full-text review will undergo data extraction (figure 1). We will use the software Covidence in the screening process and create a form to store the




Figure 1 PRISMA diagram of data extraction process. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

Table 2 Sample table to report the qualitative data from the included studies

Study	Study population	Intervention studied	Feasibility of intervention	Acceptability of intervention	Considerations for equity

data extracted in duplicate by two authors.¹⁵ Two authors will screen the studies independently and compare their findings. The authors will resolve potential disagreements through discussions, and the senior author on the team may assist as needed.

We will extract characteristic data for each eligible study, including study region/country, study year, study type, intervention exposure (dose, duration, frequency), comparison, outcomes, population characteristics detailed in subgroup analysis and risk of bias. We will also extract the information on feasibility, acceptability, equity and resource use of the intervention and report this data in separate tables (tables 2 and 3). We will extract the raw values for the number of events in the intervention and control group in case of dichotomous outcomes. To avoid reviewer bias, we are deciding a priori the order of preference for extracting outcomes when data might be available in several formats. For randomised studies, we will prefer data that require the least manipulation by authors or inference by reviewers. We will give preference denominators in the following order for mortality data: number with the definite outcome known, number randomised and child-years. For morbidity outcomes to which both survivors and non-survivors may have contributed data (eg, the incidence of diarrhoea), we will give preference to child years/months/days, number with the definite outcome known and number randomised.

For continuous outcomes from randomised trials, we will extract the raw values (eg, means and SD) rather than calculated effect sizes (eg, Cohen's d) and calculate the summary estimate ourselves. For denominators, we will preference the numbers with the definite outcome known and then number randomised. For non-randomised studies, we will prefer to extract the summary estimates (mean difference or relative risk (RR)) adjusted for confounding variables.

If a study is only available in an abstract, we will contact the authors for full text. If the full text could not be obtained from any sources, we will include the abstract if sufficient details of study design and outcomes are available. We will try to find the protocol of each included study to assess the details of the methods. If the study protocol is not publicly available, we will contact the authors for the same. Suppose the results of the randomised trial are published in more than one report. In that case, we will

consider all the publications related to that study as one study but extract the data from all relevant publications.

Assessment of risk of bias in included studies

We will evaluate the study risk of bias with the Cochrane risk of bias (ROB V.2.0) and Cochrane ROBINS-I for RCTs and non-randomised studies of interventions, respectively.^{16 17} Two authors will assess the risk of bias separately and compare their findings and reach a consensus. In the case of conflicts, a discussion will be held to resolve the conflict. A senior author will be consulted if a consensus could be reached between the two primary authors.

The risk of bias assessment according to ROB-2 and ROBINS-I is done for each outcome and not for a particular study.^{16 17} This risk of bias assessment is different from earlier practice where a study was assessed for risk of bias, and the same assessment was applied to all the outcomes extracted from that study.¹³ We will summarise the risk of bias for each domain (such as sequence generation, allocation concealment, etc) and the overall risk of bias from the study for a particular outcome. The risk of bias will be summarised as high, low or some concerns. We will present the risk of bias assessment in the form of graphs.

Data synthesis

We will report review findings both qualitatively and quantitatively. A narrative synthesis will be used to report the characteristics of all included studies and their results. A random effects meta-analysis will be employed when at least two studies possess sufficient clinical and methodological uniformity for synthesis. If quantitative synthesis is not appropriate, results will be described according to the standard guidelines of narrative synthesis.¹⁸ The software RevMan will be used for statistical analysis.¹⁹ Dichotomous outcomes will be assessed using RR effect sizes and presented with 95% CIs. The continuous outcomes will be combined to obtain a standardised mean difference and reported with a 95% CI. We will present the meta-analysis results in the form of forest plots and examine their statistical heterogeneity and inconsistency.

We will include the data from RCTs on an intention to treat analysis basis. If an RCT does not report an intention to treat analysis, we will generate our own intention to treat analysis by using the raw values extracted from the trial. Suppose an RCT does not report the raw values but

Table 3 Sample table for a description of qualitative data on resource use

Study	Intervention studies	Resources used for production	Resources used for administration of the intervention

reports the summary estimate, such as RR. In that case, we will include the summary estimate using the generic inverse variance weighting method of meta-analysis. For observational studies, we will prefer to use a summary estimate that was adjusted for confounding factors. If a non-randomised study reports an OR as the summary estimate, we will calculate the RR if the raw data are available. If the raw data are not available to calculate an RR, we will pool the OR if the large study sample size. We will do the sensitivity analysis by excluding such studies to see whether it has any detrimental effect on the final summary estimate of the pooled analysis.

We plan to pool the data separately from observational studies and randomised studies. Suppose the number of studies is small (less than 10). In that case, we may pool both the study designs in the same analysis but do a sensitivity analysis by excluding the observational studies.

We will consider the following two comparisons.

1. No routine antibiotics compared with routine antibiotics following treatment protocols in the 2013 WHO guidelines.
2. Different approaches to the use of antibiotics (eg, types of antibiotics, doses, etc) compared with the routine antibiotics following treatment protocols in the 2013 WHO guidelines.

Studies with missing data

We will record missing data during the data extraction process. We will use imputations of missing data over recording the data as missing. If data are missing for some cases or dropouts are not reported, we will contact the study authors requesting the complete data. If a study does imputations for the missing data, we will use the imputed data. If a study does not report the raw values, we will

extract the summary estimates such as RR or risk ratio. If a study does not report mean but median, we will convert the median to mean based on the Cochrane Handbook of Systematic Reviews methods.¹³ Suppose the SD is not available for a mean, and the value cannot be calculated from available data such as SE, CI or the p value. In that case, we will contact the authors for additional information. If the SD is not available from any source for a mean, we will use the SD from a similar study. If the data are presented only in graphs, we will extract the information to the best possible approximation.

Assessment of heterogeneity

We will analyse statistical heterogeneity in the pooled data using Tau,² χ^2 and I^2 statistics. We will also assess statistical heterogeneity through visual inspection of forest plots, using the χ^2 test (assessing the p value) and calculating the Tau² and I^2 statistics. We will consider it significant statistical heterogeneity when the p value is less than 0.1, the I^2 value exceeds 50%, and the inspection of forest plots shows substantial variability in the effect of the intervention. Finally, we will perform subgroup analysis to identify reasons for eligible statistical heterogeneity.

Assessment of reporting bias

We will assess publication bias of small studies using funnel plots and regression tests for funnel plot asymmetry when the meta-analysis includes at least 10 studies.

Subgroup analysis and investigation of heterogeneity

- By different types/definitions of growth failure/faltering.
- Age at presentation (newborn (0–28 days), 1–3 months, 4–6 months).

Table 4 Description of methods for grading of overall evidence

Study design	Quality of evidence*	Lower certainty score if	Higher certainty score if
Randomised trial	High	Risk of bias	Large effect
	Moderate	► 1 Serious	+1 Large
Observational study	Low	► 2 Very serious	+2 Very large
	Very low	Inconsistency	
		► 1 Serious	
		► 2 Very serious	
		Indirectness	Dose–response
		► 1 Serious	+1 evidence of a gradient
		► 2 Very serious	
		Imprecision	All plausible confounding would:
		► 1 Serious	+1 Reduce a demonstrated
		► 2 Very serious	effect
		Publication bias	
		► 1 Likely	
		► 2 Very likely	+1 Suggest a spurious effect
			when results show no effect

*In the GRADE approach, randomised controlled trials (RCTs) start as high-quality evidence and observational studies as low-quality evidence supporting estimates of intervention effects. Five factors may lead to rating down the quality of evidence, and three factors may lead to rating up. Ultimately, the quality of evidence for each outcome falls into one of four categories, from high to very low. GRADE is 'outcome centric': rating is made for each outcome, and quality may differ indeed, is likely to differ from one outcome to another within a single study and across a body of evidence.²¹

- ▶ Gestational age: preterm birth (<37 weeks) versus full-term birth (>37 weeks).
- ▶ Birth weight: low birth weight (<2500 g) versus normal birth weight (>2500 g).
- ▶ HIV exposure: studies with participants exposed to HIV versus studies with no HIV exposure.
- ▶ Presentation: participants with oedema versus participants with no oedema.
- ▶ Comorbidities: with or without comorbidities
- ▶ Nutrition: babies breast feeding or non-breastfed babies
- ▶ Location of the treatment: inpatient or outpatient/community
- ▶ Dose of antibiotics
- ▶ Duration of antibiotics: 7 days versus >7 days
- ▶ Type of antibiotics

The subgroup analyses above will be based on individual-level data. We will try to extract the information from individual studies as available. If the information is not available from the published studies, we will write to the authors to get the required data.

Sensitivity analysis

We will complete sensitivity analysis by removing studies with a high risk of bias. We will compare results from random versus fixed-effect meta-analysis models.

Rating of overall quality of evidence

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach will be used to evaluate the overall certainty of evidence using the software GRADEpro.²⁰ The GRADE approach is a comprehensive framework used to assess the overall certainty of the evidence for an outcome using study characteristics such as study design, inconsistency, indirectness of evidence, risk of bias, publication bias and imprecision estimates. We will include the GRADE assessment results in a GRADE evidence profile that contains certainty ratings, including very low, low, moderate or high based on the evidence across studies for primary outcomes (table 4).

Amendments

We will conduct literature searches, screening of titles, selection of studies, data extraction and analysis according to the plan aforementioned in the protocol. We will report any additional analysis performed in the Methods section if such occurs.

Author affiliations

¹Division of Pediatric Gastroenterology, Department of Pediatrics, SUNY Upstate Medical University, Syracuse, New York, USA

²College of Medicine, SUNY Upstate Medical University, Syracuse, New York, USA

³College of Education, University of Oregon, Eugene, Oregon, USA

⁴Health Science Library, SUNY Upstate Medical University, Syracuse, NY, USA

⁵Division of Women and Child Health, Aga Khan University, Karachi, Pakistan

⁶Division of Women and Child Health, The Aga Khan University, Karachi, Sindh, Pakistan

⁷SickKids, Toronto, Ontario, Canada

Acknowledgements We are very thankful to Allison Daniel, Jaden BENDABENDA, Kirrily De POLNAY and Zita WEISE PRINZO for their input to improve this protocol.

Contributors AI and ZAB conceptualised the study. FFC and AI wrote the first draft of the manuscript. MF contributed to method. ET-S contributed to methods. AS and OT designed the search strategy. JD, ZAB provided the supervision as experts and contributed to methods and manuscript writing.

Funding This work is funded by World Health Organization grant number 202725572. WHO also provided technical support for this work.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Aamer Imdad <http://orcid.org/0000-0002-7026-0006>

Emily Tanner-Smith <http://orcid.org/0000-0002-5313-0664>

Jai K Das <http://orcid.org/0000-0002-2966-7162>

REFERENCES

- 1 United Nations Children's Fund (UNICEF). International bank for reconstruction and Development/The world bank. levels and trends in child malnutrition: key findings of the 2020 edition of the joint child malnutrition estimates. Geneva World Health Organization; 2020.
- 2 WHO. *Guideline: updates on the management of severe acute malnutrition in infants and children*. Geneva, 2013.
- 3 Kerac M, Blencowe H, Grijalva-Eternod C, *et al*. Prevalence of wasting among under 6-month-old infants in developing countries and implications of new case definitions using who growth standards: a secondary data analysis. *Arch Dis Child* 2011;96:1008–13.
- 4 Nasrin D, Blackwelder WC, Sommerfelt H, *et al*. Pathogens associated with linear growth faltering in children with diarrhea and impact of antibiotic treatment: the global enteric multicenter study. *J Infect Dis* 2021;224:S848–55.
- 5 Brander RL, Pavlinac PB, Walson JL, *et al*. Determinants of linear growth faltering among children with moderate-to-severe diarrhea in the global enteric multicenter study. *BMC Med* 2019;17:214.
- 6 Patwari AK, Kumar S, Beard J. Undernutrition among infants less than 6 months of age: an underestimated public health problem in India. *Matern Child Nutr* 2015;11:119–26.
- 7 Prado EL, Larson LM, Cox K, *et al*. Do effects of early life interventions on linear growth correspond to effects on neurobehavioural development? A systematic review and meta-analysis. *Lancet Glob Health* 2019;7:e1398–413.
- 8 Shekhar S, Petersen FC. The dark side of antibiotics: adverse effects on the infant immune defense against infection. *Front Pediatr* 2020;8:544460.
- 9 Alcoba G, Kerac M, Breyse S, *et al*. Do children with uncomplicated severe acute malnutrition need antibiotics? A systematic review and meta-analysis. *PLoS One* 2013;8:e53184.
- 10 Stam J, van Stuijvenberg M, Grüber C, *et al*. Antibiotic use in infants in the first year of life in five European countries. *Acta Paediatr* 2012;101:929–34.
- 11 Droste JH, Wieringa MH, Weyler JJ, *et al*. Does the use of antibiotics in early childhood increase the risk of asthma and allergic disease? *Clin Exp Allergy* 2000;30:1548–53.

- 12 Trasande L, Blustein J, Liu M, *et al.* Infant antibiotic exposures and early-life body mass. *Int J Obes* 2013;37:16–23.
- 13 Higgins JPT TJ, Chandler J, Cumpston M, eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.2*, 2021. www.training.cochrane.org/handbook
- 14 Page MJ, McKenzie JE, Bossuyt PM, *et al.* The prisma 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- 15 Covidence systematic review software. Veritas health innovation. Melbourne, Australia; 2021. www.covidence.org
- 16 Sterne JAC, Savović J, Page MJ, *et al.* Rob 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:l4898.
- 17 Sterne JA, Hernán MA, Reeves BC, *et al.* ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:i4919.
- 18 Campbell M, McKenzie JE, Sowden A, *et al.* Synthesis without meta-analysis (swim) in systematic reviews: reporting guideline. *BMJ* 2020;368:l6890.
- 19 The Nordic Cochrane Center TCC. *Review manager (RevMan)*. 5.3. Copenhagen, 2014.
- 20 Evidence Prime, Inc. GRADEpro GDT: GRADEpro Guideline development tool. McMaster University; 2020.
- 21 Guyatt G, Oxman AD, Akl EA, *et al.* Grade guidelines: 1. introduction-grade evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383–94.

Appendix 1

Search Strategies for Growth Failure AND Antibiotics

PubMed

(("Infant"[MeSH Terms] OR "Child"[MeSH Terms:noexp] OR "Premature Birth"[MeSH Terms] OR
 ("infant*"[Title/Abstract] OR "infancy"[Title/Abstract] OR
 "baby"[Title/Abstract] OR
 "babies"[Title/Abstract] OR "neonat*"[Title/Abstract] OR "neo
 nat*"[Title/Abstract] OR
 "newborn*"[Title/Abstract] OR "new born*"[Title/Abstract] OR "newly
 born*"[Title/Abstract] OR
 "preterm"[Title/Abstract] OR "preterms"[Title/Abstract] OR "pre
 term"[Title/Abstract] OR "pre
 terms"[Title/Abstract] OR "lbw"[Title/Abstract] OR "vlbw"[Title/Abstract] OR
 "elbw"[Title/Abstract]
 OR "child*"[Title/Abstract] OR "youth"[Title/Abstract] OR
 "juvenile*"[Title/Abstract]) OR
 ("premature*"[Title/Abstract] AND ("birth*"[Title/Abstract] OR
 "born"[Title/Abstract] OR
 "deliver*"[Title/Abstract])) OR ("low"[Title/Abstract] AND
 ("birthweight*"[Title/Abstract] OR "birth
 weight*"[Title/Abstract])) AND ("Malnutrition"[MeSH Terms] OR "Failure to
 Thrive"[MeSH Terms]
 OR "Growth Disorders"[MeSH Terms:noexp] OR "Severe Acute Malnutrition"[MeSH
 Terms] OR
 "Wasting Syndrome"[MeSH Terms:noexp] OR "Infant Nutrition Disorders"[MeSH
 Terms:noexp]
 OR ("growth failure*"[Title/Abstract] OR "growth falter*"[Title/Abstract] OR "slow
 growth"[Title/Abstract] OR "slowed growth"[Title/Abstract] OR
 "Malnutrition"[Title/Abstract] OR
 "malnourish*"[Title/Abstract] OR "Severe Acute Malnutrition"[Title/Abstract] OR
 "SAM"[Title/Abstract] OR "wasting"[Title/Abstract] OR
 "wasted"[Title/Abstract] OR "growth
 restrict*"[Title/Abstract] OR "growth retard*"[Title/Abstract] OR "Failure to
 Thrive"[Title/Abstract]
 OR "FTT"[Title/Abstract] OR "growth disorder*"[Title/Abstract] OR "growth
 arrest*"[Title/Abstract]
 OR "growth deficien*"[Title/Abstract] OR "growth disturb*"[Title/Abstract])) AND
 ("Anti-Bacterial
 Agents"[Pharmacological Action] OR "Anti-Bacterial Agents"[MeSH Terms] OR
 ("antibiotic*"[Title/Abstract] OR "anti bacterial agent*"[Title/Abstract] OR
 "antibacterial*"[Title/Abstract] OR "anti bacterial*"[Title/Abstract] OR
 "bactericid*"[Title/Abstract]
 OR "bacteriocid*"[Title/Abstract] OR "anti mycobacterial*"[Title/Abstract] OR
 "antimycobacterial*"[Title/Abstract])) NOT ("Animals"[MeSH Terms] NOT
 ("Animals"[MeSH
 Terms] AND "Humans"[MeSH Terms]))

CINAHL

S1 (MH "Infant+")

S2 (MH "Child")

S3 (MH "Childbirth, Premature")

S4 TI (infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new

born*" OR "newly born*" OR preterm OR preterms OR "pre term*" OR "pre terms*" OR lbw OR vlbw
 OR elbw OR child* OR youth OR juvenile*)
 S5 AB (infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term*" OR "pre terms*" OR lbw OR vlbw
 OR elbw OR child* OR youth OR juvenile*)
 S6 TI (premature* N2 (birth* OR born OR deliver*))
 S7 AB (premature* N2 (birth* OR born OR deliver*))
 S8 TI (low N2 (birthweight* OR "birth weight*"))
 S9 AB (low N2 (birthweight* OR "birth weight*"))
 S10 S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9

S11 (MH "Malnutrition")
 S12 (MH "Failure to Thrive")
 S13 (MH "Growth Disorders")
 S14 (MH "Wasting Syndrome")
 S15 (MH "Infant Nutrition Disorders")
 S16 TI ((grow*) N2 (fail* OR falter* OR slow* OR restrict* OR retard* OR deficien* OR arrest* OR disturb* OR disorder*))
 S17 AB ((grow*) N2 (fail* OR falter* OR slow* OR restrict* OR retard* OR deficien* OR arrest* OR disturb* OR disorder*))
 S18 TI ((fail*) W2 (thrive))
 S19 AB ((fail*) W2 (thrive))
 S20 TI (Malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR SAM OR wasting OR wasted OR FTT)
 S21 AB (Malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR SAM OR wasting OR wasted OR FTT)
 S22 S11 OR S12 OR S13 OR S14 OR S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S21
 S23 (MH "Antibiotics+")
 S24 TI (antibiotic* OR "anti bacterial*" OR "antibacterial*" OR bactericid*OR bacteriocid* OR "anti mycobacterial*" OR "antimycobacterial*")
 S25 AB (antibiotic* OR "anti bacterial*" OR "antibacterial*" OR bactericid* OR bacteriocid* OR "anti mycobacterial*" OR "antimycobacterial*")
 S26 S23 OR S24 OR S25
 S27 (MH "Animals") NOT ((MH "Animals") AND (MH "Human"))
 S28 S10 AND S22 AND S26
 S29 S28 NOT S27
 S30 S28 NOT S27 Exclude Medline Records

Embase
 #1'infant'/exp OR 'newborn'/exp OR 'prematurity'/exp OR 'low birth weight'/exp OR 'very low birth weight'/exp OR 'extremely low birth weight'/exp OR 'juvenile'/de OR 'child'/de
 #2 infant*:ti,ab OR infancy:ti,ab OR baby:ti,ab OR babies:ti,ab OR neonat*:ti,ab OR 'neo nat*':ti,ab OR newborn*:ti,ab OR 'new born*':ti,ab OR 'newly born*':ti,ab OR preterm:ti,ab OR preterms:ti,ab
 OR 'pre term':ti,ab OR 'pre terms':ti,ab OR lbw:ti,ab OR vlbw:ti,ab OR elbw:ti,ab OR child*:ti,ab OR youth:ti,ab OR juvenile*:ti,ab
 #3 (premature* NEAR/2 (birth* OR born OR deliver*)):ti,ab

#4 (low NEAR/2 (birthweight* OR 'birth weight*')):ti,ab

#5 #1 OR #2 OR #3 OR #4

#6 'malnutrition'/exp OR 'failure to thrive'/exp OR 'growth disorder'/de OR
'wasting syndrome'/de
OR 'growth retardation'/de

#7(grow* NEAR/2 (fail* OR falter* OR slow* OR restrict* OR retard* OR deficien* OR arrest* OR
disturb* OR disorder*)):ti,ab

#8 (fail* NEXT/2 thrive):ti,ab

#9 malnutrition:ti,ab OR malnourish*:ti,ab OR 'severe acute malnutrition';ti,ab OR sam:ti,ab OR
wasting:ti,ab OR wasted:ti,ab OR ftt:ti,ab

#10 #6 OR #7 OR #8 OR #9

#11'antibiotic agent'/exp OR 'bactericide'/de OR 'antimycobacterial agent'/de

#12 antibiotic*:ti,ab OR 'anti bacterial*';ti,ab OR 'antibacterial*';ti,ab OR bactericid*:ti,ab OR
bacteriocid*:ti,ab OR antimycobacterial*:ti,ab OR 'anti mycobacterial*';ti,ab

#13 #11 OR #12

#14 #5 AND #10 AND #13

#15 #14 NOT ('animals'/exp NOT 'humans'/exp)

#16 #15 NOT [medline]/lim

CENTRAL

#1 MeSH descriptor: [Infant] explode all trees

#2 MeSH descriptor: [Child] this term only

#3 MeSH descriptor: [Premature Birth] explode all trees

#4 infant*:ti,ab OR infancy:ti,ab OR baby:ti,ab OR babies:ti,ab OR neonat*:ti,ab OR newborn*:ti,ab
OR preterm:ti,ab OR preterms:ti,ab OR lbw:ti,ab OR vlbw:ti,ab OR elbw:ti,ab OR child*:ti,ab OR
youth:ti,ab OR juvenile*:ti,ab

#5 (neo NEXT nat* OR new NEXT born* OR newly NEXT born* OR pre NEXT term OR pre NEXT
terms):ti,ab

#6 (premature* NEAR/2 (birth* OR born OR deliver*)):ti,ab

#7 (low NEAR/2 (birthweight* OR birth NEXT weight*)):ti,ab

#8 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7

#9 MeSH descriptor: [Malnutrition] explode all trees

#10 MeSH descriptor: [Failure to Thrive] explode all trees

#11 MeSH descriptor: [Growth Disorders] this term only

#12 MeSH descriptor: [Severe Acute Malnutrition] explode all trees

#13 MeSH descriptor: [Wasting Syndrome] this term only

#14 MeSH descriptor: [Infant Nutrition Disorders] this term only

#15 ((grow*) NEAR/2 (fail* OR falter* OR slow* OR restrict* OR retard* OR deficien* OR arrest* OR
disturb* OR disorder*)):ti,ab

#16 ((fail*) NEXT/2 (thrive)):ti,ab

#17 (severe NEXT acute NEXT malnutrition):ti,ab

#18 malnutrition:ti,ab OR malnourish*:ti,ab OR sam:ti,ab OR wasting:ti,ab OR wasted:ti,ab OR
ftt:ti,ab

#19 #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18

#20 MeSH descriptor: [Anti-Bacterial Agents] explode all trees

#21 antibiotic*:ti,ab OR antibacterial*:ti,ab OR bactericid*:ti,ab OR bacteriocid*:ti,ab OR
antimycobacterial*:ti,ab

#22 (anti NEXT bacterial*):ti,ab

#23 (anti NEXT mycobacterial*):ti,ab

#24 #20 OR #21 OR #22 OR #23

#25 #8 AND #19 AND #24

#26 MeSH descriptor: [Animals] explode all trees

#27 MeSH descriptor: [Humans] explode all trees

#28 (#26 NOT (#26 AND #27))

#29 #25 NOT #28

#30 "accession number" NEAR pubmed

#31 #29 NOT #30

Scopus

((TITLE-ABS (infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term" OR

"pre terms" OR lbw OR vlbw OR elbw OR child* OR youth OR juvenile*)) OR (TITLE-ABS (low W/2 (birthweight* OR "birth weight*"))) OR (TITLE-ABS (premature* W/2 (birth* OR born OR deliver*)))) AND ((TITLE-ABS ((grow*) W/2 (fail* OR falter* OR slow* OR restrict* OR retard* OR deficien* OR arrest* OR disturb* OR disorder*))) OR (TITLE-ABS ((fail*) PRE/2 (thrive))) OR (TITLE-ABS (malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR sam OR wasting OR wasted OR ftt))) AND (TITLE-ABS (antibiotic* OR "anti bacterial*" OR "antibacterial*" OR bactericid* OR bacteriocid* OR "anti mycobacterial*" OR "antimycobacterial*")) AND NOT INDEX (medline)

Web of Science

#1 TI=(infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term" OR "pre terms" OR

lbw OR vlbw OR elbw OR child* OR youth OR juvenile*)

#2 AB=(infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term" OR "pre terms" OR

lbw OR vlbw OR elbw OR child* OR youth OR juvenile*)

#3 TI=(premature* NEAR/2 (birth* OR born OR deliver*))

#4 AB=(premature* NEAR/2 (birth* OR born OR deliver*))

#5 TI=(low NEAR/2 (birthweight* OR "birth weight*"))

#6 AB=(low NEAR/2 (birthweight* OR "birth weight*"))

#7 #6 OR #5 OR #4 OR #3 OR #2 OR #1

#8 TI=((grow*) NEAR/2 (fail* OR falter* OR slow* OR restrict* OR retard* OR deficien* OR arrest* OR disturb* OR disorder*))

#9 AB=((grow*) NEAR/2 (fail* OR falter* OR slow* OR restrict* OR retard* OR deficien* OR arrest* OR disturb* OR disorder*))

#10 TI=((fail*) NEAR/2 (thrive))

#11 AB=((fail*) NEAR/2 (thrive))

#12 TI=(Malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR SAM OR wasting OR wasted OR FTT)

#13 AB=(Malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR SAM OR wasting OR wasted OR FTT)

#14 #13 OR #12 OR #11 OR #10 OR #9 OR #8

#15 TI=(antibiotic* OR "anti bacterial*" OR "antibacterial*" OR bactericid* OR bacteriocid* OR "anti mycobacterial*" OR "antimycobacterial*")

#16 AB=(antibiotic* OR "anti bacterial*" OR "antibacterial*" OR bactericid* OR bacteriocid* OR "anti mycobacterial*" OR "antimycobacterial*")

"anti mycobacterial*" OR "antimycobacterial*")

#17 #16 OR #15

#18 #17 AND #14 AND #7

LILACS

(mh:(infant)) OR ((mh:(child))) OR ((mh:("Premature Birth"))) OR ((ti:((premature* AND (birth* OR born OR deliver*)))) OR ((ab:((premature* AND (birth* OR born OR deliver*)))) OR ((ti:((low AND (birthweight* OR "birth weight*")))) OR ((ab:((low AND (birthweight* OR "birth weight*")))) OR
 ((ti:(infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term*" OR "pre terms*" OR lbw OR vlbw OR elbw OR child* OR youth OR juvenile*))) OR ((ab:(infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term*" OR "pre terms*" OR lbw OR vlbw OR elbw OR child* OR youth OR juvenile*))) AND
 ((mh:(malnutrition)) OR ((mh:("Failure to Thrive"))) OR ((mh:("Growth Disorders"))) OR ((mh:("Severe Acute Malnutrition"))) OR ((mh:("Wasting Syndrome"))) OR ((mh:("Infant Nutrition Disorders"))) OR ((ti:("growth failure*" OR "growth falter*" OR "slow growth*" OR "slowed growth*" OR malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR sam OR wasting OR wasted OR "growth restrict*" OR "growth retard*" OR "Failure to Thrive" OR ftt OR "growth disorder*" OR "growth arrest*" OR "growth deficien*" OR "growth disturb*"))) OR ((ab:("growth failure*" OR "growth falter*" OR "slow growth*" OR "slowed growth*" OR malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR sam OR wasting OR wasted OR "growth restrict*" OR "growth retard*" OR "Failure to Thrive" OR ftt OR "growth disorder*" OR "growth arrest*" OR "growth deficien*" OR "growth disturb*")))) AND ((mh:("Anti-Bacterial Agents")) OR ((ti:(antibiotic* OR "anti bacterial agent*" OR antibacterial* OR "anti bacterial*" OR "bactericid*" OR "bacteriocid*" OR "anti mycobacterial*" OR antimycobacterial*))) OR ((ab:(antibiotic* OR "anti bacterial agent*" OR antibacterial* OR "anti bacterial*" OR "bactericid*" OR "bacteriocid*" OR "anti mycobacterial*" OR antimycobacterial*)))) AND NOT ((mh:(animals)) AND NOT ((mh:(animals)) AND ((mh:(humans)))))
 Global Index Medicus
 (mh:(infant)) OR ((mh:(child))) OR ((mh:("Premature Birth"))) OR ((ti:((premature* AND (birth* OR born OR deliver*)))) OR ((ab:((premature* AND (birth* OR born OR deliver*)))) OR ((ti:((low AND (birthweight* OR "birth weight*")))) OR ((ab:((low AND (birthweight* OR "birth weight*")))) OR
 ((ti:(infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term*" OR "pre terms*" OR lbw OR vlbw OR elbw OR child* OR youth OR juvenile*))) OR ((ab:(infant* OR infancy OR baby OR babies OR neonat* OR "neo nat*" OR newborn* OR "new born*" OR "newly born*" OR preterm OR preterms OR "pre term*" OR "pre terms*" OR lbw OR vlbw OR elbw OR child* OR youth OR juvenile*))) AND
 ((mh:(malnutrition)) OR ((mh:("Failure to Thrive"))) OR ((mh:("Growth Disorders"))) OR

((mh:("Severe Acute Malnutrition"))) OR ((mh:("Wasting Syndrome"))) OR
 ((mh:("Infant Nutrition
 Disorders"))) OR ((ti:("growth failure*" OR "growth falter*" OR "slow
 growth" OR "slowed growth"
 OR malnutrition OR malnourish* OR "Severe Acute Malnutrition" OR sam OR wasting OR wasted
 OR "growth restrict*" OR "growth retard*" OR "Failure to Thrive" OR ftt OR
 "growth disorder*" OR
 "growth arrest*" OR "growth deficien*" OR "growth disturb*"))) OR
 ((ab:("growth failure*" OR
 "growth falter*" OR "slow growth" OR "slowed growth" OR malnutrition OR
 malnourish* OR
 "Severe Acute Malnutrition" OR sam OR wasting OR wasted OR "growth restrict*" OR
 "growth
 retard*" OR "Failure to Thrive" OR ftt OR "growth disorder*" OR "growth
 arrest*" OR "growth
 deficien*" OR "growth disturb*")))) AND ((mh:("Anti-Bacterial Agents")) OR
 ((ti:(antibiotic* OR
 "anti bacterial agent*" OR antibacterial* OR "anti bacterial*" OR "bactericid*"
 OR "bacteriocid*"
 OR "anti mycobacterial*" OR antimycobacterial*))) OR ((ab:(antibiotic* OR "anti bacterial
 agent*"
 OR antibacterial* OR "anti bacterial*" OR "bactericid*" OR "bacteriocid*" OR
 "anti mycobacterial*"
 OR antimycobacterial*)))) AND NOT ((mh:(animals)) AND NOT ((mh:(animals)) AND
 ((mh:(humans)))) AND (collection_gim:("IMSEAR" OR "WPRIM" OR "IMEMR"
 OR "AIM"))