| Outcome | Screen positive; Aspirin Group 3A N=13 | Screen negative; No Aspirin Group 3B N=171 | OR (95% CI) |
|---|--|--|------------------|
| Preeclampsia No. (%) | 2 (15.4) | 6 (3.5) | 5.0 (0.9 – 27.7) |
| Pre-eclampsia <34- | 0 (0) | 2 (1.2) | - |
| weeks | | | |
| Pre-eclampsia <37- weeks | 2 (15.4) | 2 (1.2) | 15.4 (2.0 – 120) |
| Birthweight <10 th centile No. (%) | 4 (30.7) | 21 (12.3) | 3.2 (0.9 – 11.2) |
| Pre-term delivery <34 weeks No. (%) | 1 (7.7) | 1 (0.6) | 14.1 (0.8 – 240) |
| NICU admission No. | 0 (0) | 9 (5.3) | - |
| Outcome No. (%) | | | |
| Alive at 6-weeks | 13 (100) | 169 (98.8) | |
| Stillbirth | 0 (0) | 2 (1.2) | |
| Neonatal death | 0 (0) | 0 (0) | |

Table S2 - Secondary outcome measures in Group 3 (screen and treat) $\,$

(Expressed as average and standard deviation unless otherwise stated)

Note: ORs are not presented when number of events is 0 in the Screen –positive group.